



Nutritional Impact of Housefly Maggot Meal on Growth of African Catfish (*Clarias Gariepinus*)

Nzegwu Johnson Onyebuchi ^a, Egboka Tochukwu Praise ^b,
Anukwuorji Chidozie Azubuike ^{b*}
and Egboka Chidiebele Roseline ^b

^a Department of Zoology, Faculty of Bioscience, Nnamdi Azikiwe University, Awka, Nigeria.

^b Department of Botany, Faculty of Bioscience, Nnamdi Azikiwe University, Awka, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

<https://prh.ikpress.org/review-history/12533>

Original Research Article

Received: 13/09/2024

Accepted: 16/11/2024

Published: 28/11/2024

ABSTRACT

In an attempt to successfully replace fish meal with Maggot meal in the diet of *Clarias gariepinus* for optimal growth and nutrient utilization, a 49 days (7 weeks) laboratory feeding trial was conducted to evaluate the growth of *Clarias gariepinus* fingerlings with the average initial weight of 5.36 ± 0.96 g and average initial length of 4.20 ± 0.05 cm, using housefly larva (*Musca domestica*) Maggot meal from cow dung as the protein source in place of fish meal in their diet. A total of 60 fingerlings were used. They were stocked 10 fingerlings in a 70 litre bowl. Each treatment was replicated twice. Treatment 1 (0% maggot inclusion), Treatment 2 (50% maggot inclusion) and Treatment 3 (53% maggot inclusion). At the end of the 49 days (7 weeks), the fingerlings fed with 53% maggot inclusion recorded the highest mean weight (7.93g) and mean length (12.01cm). The data obtained

*Corresponding author: E-mail: dozygreat@gmail.com;

Cite as: Onyebuchi, Nzegwu Johnson, Egboka Tochukwu Praise, Anukwuorji Chidozie Azubuike, and Egboka Chidiebele Roseline. 2024. "Nutritional Impact of Housefly Maggot Meal on Growth of African Catfish (*Clarias Gariepinus*)". *Asian Journal of Research in Biology* 7 (1):36-43. <https://journalbiology.org/index.php/AJRiB/article/view/35>.

were subjected to one-way analysis of variance (ANOVA) while Fisher's least significant difference test ($P \leq 0.05$) was used to compare treatment means. Treatment 1 fish meal inclusion has the highest specific growth rate (12.9) but not significantly different and 53% maggot inclusion has the highest PER (1.63) but the least value was recorded in fish meal diet. The highest value for FCR was recorded in 53% maggot inclusion (0.83) while treatment 1 has the highest SGR and treatment 2 containing 50% maggot meal inclusion has the least survival rate while treatment 1 and 3 has the highest survival rate. There was no significant different ($P > 0.05$) in all the parameters. Conclusively maggot meal can be used successfully in place of fish meal in the diet of *Clarias gariepinus*.

Keywords: Maggot-meal; treatment; inclusion; protein; proximate.

1. INTRODUCTION

"The increasing cost of feed has been at alarming rate and this has been affecting the development and expansion of aquaculture in African countries particularly Nigeria" [1]. "Fish meal been the major protein source in the fish diet constitutes the highest cost thereby making the price of the feed to rise exponentially. Therefore, several attempts have been made to substitute fish meal with other animal protein sources such as earth worm, shrimp waste, insect and plant protein sources such as sun flower, rape seed, soy bean meal and cottonseed meal. However, they cannot replace fish meal wholly but partially due to the presence of chitin in their exoskeleton. In the same vein problem of anti- nutritional factor in tropical legumes have limited their usage and direct incorporation into animal feeds" [2]. "Maggot meal is an animal protein source produced from waste, it has been reported to be highly nutritive with crude protein ranging between 43.9 and 62.4%, lipid 12.5 and 21%, and crude fibre 5.8 and 8.2%" [3]. "Maggot meal is also rich in phosphorus, trace elements and B complex. The cost of harvesting and processing one kilogram of maggot meal is smaller compared to the cost of 1kg of fish meal, thereby showing the cost effectiveness of using maggot meal in the diet of African catfish. It is on record that the replacement of fish meal with 50% and 100% maggot meal reduces the production cost per kilogram by approximately 18 to 28% due to the low cost of this ingredients" [3]. "Presently the exorbitant market prices of conventional feed materials such as fish meal and soybean meal necessitated the need for the development of other protein sources for aquaculture as a new area of research" [4]. "The increasing cost of fish feed has been at an alarming rate and this has affected the development and expansion of aquaculture in African countries" [5]. "The need for more research for vital protein augments to make

affordable fishmeal and thus increase the production of catfish becomes eminent" [6]. "The research for suitable and cost-effective alternative protein sources for use in industrial aqua feeds will be the most critical factor in the development of intensive aquaculture" [1]. "Insect meals are nutritious and healthy alternatives to fishmeal because of its rich nutritional values especially protein, fat and minerals" [7]. "Alternative protein sources can provide solutions to problems faced by small-scale farmers such as high costs of production and the resultant loss in revenue due to low farm-gate prices" [4].

There are myriads of reports on the use of other sources of protein other than the conventional sources as alternative protein in fish feed. However, there is a dearth of information on the use of housefly maggots as alternative. It is against the background of the above scenario that this research was conducted to ascertain the effects of housefly maggot on the growth of African catfish. This study therefore evaluates the nutritive value and growth performance of *Clarias gariepinus* fed with maggot meal.

2. MATERIALS AND METHODS

2.1 Experimental Fish Collection

The experimental fish, fingerlings of the catfish *Clarias gariepinus* were collected from Ejiamatu fish farm, Aguoye in Anambra State. The weight of the samples collected range between 5.36g. A total of sixty (60) fish were collected. The fish were transported to the Zoology hatchery unit in a plastic bucket with well oxygenated water. The fish samples were acclimated to laboratory conditions for seven days in big plastic tanks of 70 liters volume.

2.2 Experimental Design

Six (6) bowls of 70litres volume were used. The bowls were of varying colors (blue and green) comprising of three (3) treatments and their

replicates respectively. The experimental design uses a completely randomized design with 3 treatments with 3 replications (10 fish in each replication). The treatments given are T0, T1, T2 and T3. The experimental design is shown as follows: Compounded feed consisting of fishmeal, compounded feed containing 50% of maggot meal and compounded feed containing 53% of maggot meal.

The fish were stocked at the density of ten (10) fingerlings per plastic bowl and covered with nets to prevent the fish from jumping out. During the period of Acclimatization, the fish were fed Coppens at 5% body weight twice daily (Okoye et al., 2001) with the experimental diets of 40% crude protein. The main source of water used was well water located at the hatchery unit. The fish in each tank were selected at random and were fed at 5% of their body weight with the experimental diets. The fish were fed twice daily between 8.00am and 5.00pm.

2.3 Weighing and Measurement

All the fish in each group of feeding regime were weighed individually using water displacement method on an electronic sensitive scale (Labtech brand, model BL7501 made by LAB-TECH India) at the beginning of the experiment and thereafter on a weekly basis to determine the growth rate of the fish. The water in each treatment was sieved using a sieve and the fingerlings were weighed individually using an electronic sensitive meter which gives the accurate weight of the fingerling when placed on the meter's plate-like surface after which they were measured using a meter rule and the length and weight were recorded respectively.

2.4 Management of Experimental Systems

Changing of water in all the treatments was done twice a week.

The following water quality parameters were measured carefully at the beginning and at the end of the experiment: Temperature, pH, Dissolved oxygen.

- Temperature: This was taken at the beginning and at the end of the experiment using mercury in-glass thermometer.
- pH: pH was measured using pH meter.

2.5 Maggot Production

A 25 kg of fresh cattle dung, 5 kg of spent grain and saw dust were mixed together to a thickness of 5 cm to constitute substrate. The mixture was spread in a box of 1.4 m by 1.4 m screened with plastic net with one end open. The dung product were left to ferment for 48 h and the odour released attracted flies to perch on the mixture, then the open side was closed the second day. The eggs laid by the female flies on the substrate hatched into larvae within two days, after which the larval matured and was harvested. The whole process took five days. The larvae were harvested and oven dried to a constant weight, after which the dried maggots were milled into a meal and kept in an air tight container at room temperature.

2.6 Experimental Diets

Feed ingredients were purchased from a feed ingredient store in a container of plastics. The major feed ingredients used were yellow maize, wheat bran, groundnut cake (GNC), fishmeal, vitamins, mineral premix, oyster shell three isonitrogenous diets of 40% crude protein were formulated: Treatment 1 (Trt 1) contained 0% of maggot meal (control), Trt 2 (50% maggot meal, Trt 3 (53.% maggot meal), with these ratios of maggot meal to fish meal 0, 1:1, 1:2 respectively. The feed were made into pellets with the use of pelletizing machine; pellets were sun dried for three days to reduce the moisture content and to prevent deterioration. The feeds were packed and stored in air tight container at room temperature

2.7 Proximate Analysis of the On-Farm Feed

The proximate analysis of the on-farm feed was carried out at Springboard Research Laboratory located at Udoka Estate, Awka in Anambra state.

Experimental diet (On-farm feed) was analyzed for proximate composition according to AOAC [8].

2.8 Determination of Moisture Content

Moisture content in feed sample was calculated by using Association of the Official Analytical Chemists [8] procedure via the sample drying method in an oven at 135°C temperatures for 2 hours till constant weight. The moisture content was calculated as follows:

Moisture content (%) = Weight of dried sample / original wet weight x100

2.9 Determination of Ash Content

Ash content in feed samples was calculated by using Association of the Official Analytical Chemists [8] method at 600°C for 2 hours. Percentage of ash content was measured by the resulting inorganic residue. Weight of ash obtained in percentage as follows:

Ash (%) = Weight of ash / original wet weight x100

Crude Fiber (%) = Loss in weight / Weight of sample x100

2.10 Determination of Crude Fat

The crude fat from the feed sample was extracted by Soxhlet using Association of the Official Analytical Chemists [8] methods for 4 hours at condensation rate of 5-6 drops/ second, dry extract for 30 minutes at 100 °C cool. Crude protein contents was determined gravimetrically and stored at 4 C until analysis.

2.11 Nitrogen Free Extract

NFE was determined by subtracting % crude protein, % crude fibre, %moisture, %Ash and %Ether extract from 100 using the formula:

NFE = 100 - (%moisture ± % crude protein ± % Ether extract ± %crude fibre ± % Ash).

2.12 Metabolizable Energy

Metabolizable energy was calculated based on the standard physiological values of 4.5 kcal/g protein, 3.3 kcal/g carbohydrates and 8 kcal/g fat [9].

2.13 Growth and Food Evaluation Data

i). Specific Growth Rate

SGR% = $\frac{\ln W_2 - \ln W_1}{T-t} \times 100$
Where;

W₁, is the initial weight (gram at time t).
W₂, is the final weight (gram at time T)
t= initial time

T= final time

The food conversion ratio was calculated thus:

ii). F.C.R = Feed intake / Fish weight gain

The study lasted for a period of seven weeks.

iii. Protein Efficiency Ratio (P.E.R):

This was determined using the formula;

PER = Weight gain / Weight of Protein intake x100

Percentage survival rate:

Percentage survival rate = $\frac{\text{Final number of fish}}{\text{Initial number of fish}} \times 100$

2.14 Statistical Analysis of Data

Statistical analysis was carried out using GenStat Release 12.1 Software. The data were subjected to one-way analysis of variance (ANOVA). Fisher's least significant difference test ($P \leq 0.05$) was used to compare treatment means.

3. RESULTS

From the result presented in the Fig. 1, it can be deduced that there is a successive increase in the weight of the experimental fish fed different inclusion of maggot meal than those obtain from fishmeal diet. These data is higher than the result obtained before the experimental research. But from the statistical analysis, there was no significant different among the experimental groups fed the two diets (fishmeal and maggot meal) ($P>0.05$).

From the result presented in Fig. 2 of, it can be deduced that there was a recorded increase in the weekly mean length among the experimental group but the result was subjected to statistical analysis which show that there was no significant difference among the experimental group as ($P>0.05$).

From the experimental result presented above on Table 2, treatment 3 consumed more feed made of maggot meal, followed by treatment 1 that was fed with fishmeal diet. Treatment 2 has the least food consumption and as a result has the least food conversion ratio. From the result presented above also, it can be deduced that the feed

made with maggot meal has the highest protein efficiency treatment 1 seems to have a high SGR but from the statistical analysis, it has deduced that there was no significant difference in the growth indices of the fish feed the experimental diets as ($P>0.05$).

Table 3 above shows the result of the percentage survival rate of *Clarias gariepinus* fingerlings fed on two different diets (fishmeal and maggot meal) from the table, it can be deduced that *Clarias gariepinus* fingerlings fed on fishmeal feed and 53% maggot meal inclusion had the highest percentage survival rate (75%). Those fed 50% maggot meal feed has the least percentage survival rate (70 %.) There is no significant difference ($P>0.05$).

4. DISCUSSION

The findings of this study showed the prospects of using maggot meal as alternative to fishmeal. The proximate composition of the experimental diet as presented in Table 1 depicted the composition of the feed with respect to different

protein contents at varying levels. The conventional fish meal has been proven to be rich in essential nutrients needed for fish growth and development. In this study, maggot meal was well utilized by *C. gariepinus* and this result in good fish performance as observed in the various parameters measured (weight and length). This corroborates with the finding of several researchers who postulates that maggot meal, like other animal protein sources, is well accepted and utilized by the fish [10,11].

The increase in the growth of fish fed all the experimental diets could be as a result of the rich nutrient contents of the various experimental diets. However, it's pertinent to note that no statistically significant was observed in the weight of the fish across the experimental diets used. The non-significant difference of the evaluated parameters of growth and nutrient utilization among the three (3) experimental treatments points to the fact that maggot meal can successfully be used as alternative to fish meal in fish diets. A review of extant literatures

Table 1. Proximate composition of experimental diets

Parameters	Treatment 1	Treatment 2	Treatment 3
CP	44.20	43.50	44.23
CF	3.74	3.76	3.74
EE	4.76	4.60	4.50
ASH	13.37	12.77	13.11
DM	9.95	8.90	9.90
NFE	23.98	25.47	24.52


CP=Crude protein, CF=Crude fatty acid, EE-Crude fiber, ASH=Ash, DM=Moisture, NFE=Nitrogen free extract.

Table 2. Specific growth rate (SGR), protein efficiency ratio (PER) and food conversion ratio (FCR) of *Clarias gariepinus* fingerlings feed on different combinations of fish meal and maggot feeds for seven (7 weeks)

Growth indices	Treatment 1	Treatment 2	Treatment 3
Food conversion ratio	0.82±0.05	0.73±0.01	0.83±0.01
Protein efficiency Ratio	1.52±0.01	1.56±0.02	1.63±0.02
Specific growth rate	12.94±2.34	10.87±0.72	9.76±0.14

Table 3. Percentage survival rate of *Clarias gariepinus* fingerlings feed on two experimental feed (fishmeal and maggot meal for 49days)

Growth indices	Treatment 1	Treatment 2	Treatment 3
No of fish in each replicates	10	10	10
Replicates			
No of fish that survived	7.50	7.00	7.50
Survival rate (%)	75	70	75



Key: T₁ = 0% maggot meal; T₂ = 50% maggot meal; T₃ = 53% maggot meal

Fig. 1. Weekly mean weight increase and analysis of variance of *Clarias gariepinus* fingerlings fed on different combination of fishmeal diet and maggot meal feeds for seven (7 weeks) is presented below




Fig. 2. Weekly mean length increase of *Clarias gariepinus* fed with varied inclusion of maggot meal

and the documentations of some researchers suggests that maggot can improve the growth of catfish (*C. gariepinus*), [12,13,1,5]. The key findings of this research reinforces the reports of so many researchers on the importance of animal protein such as maggot meal in growth, development and nutrient utilization parameters of fish [10]. It was also documented by Ekeocha et al. (2023) that the biological value of protein is

close to those of groundnut cake and soya bean meal, thus maggot combination with them as alternative to fish meal is reasonable. This also agrees with the findings of Ogunji et al. [11] who reported that the strong growth and nutrient utilization capacity of fish fed a maggot meal supplemented diet stems from its high nutrient composition. Fasakin et al., [14] attributed the reduction in growth performance of experimental

fish fed full- fat maggot to low protein digestibility of the feed stuff among other reasons. Ajani et al., [3] reported maggot meal crude protein content range of 40% to 61.4% according to them, the value could be attributed to the Processing, drying or storage method as well as the quality of poultry droppings used to produce maggot. More work will be needed to standardize the method of maggot meal production. The rich nutritional content of maggot meal makes it better than other alternative protein source especially plant protein. The non-significant difference in the value of food conversion ratio of the treatment diets is a possibly indicative that both protein source compared favourably in the feed to flesh conversion. It has been reported that the biological value of maggot meal is equivalent to that of whole fish meal [3]. This fact is in accordance to the result obtained in the present study where protein efficiency ratio values were not significantly different. The percentage survival showed that there were mortality recorded during the period of study and this might be as a result of stress, water quality differences. The utilization of maggot meal will thus pave way for cheaper and nutritionally rich aqua feeds [15].

5. CONCLUSION AND RECOMMENDATION

The findings of this study revealed that maggot meal is an alternative to the conventional fish meal and other protein sources used in the production of fish. Maggot meal is relatively cheaper and readily available, hence its utilization will no doubt lower the cost of fish feeds and by extension the price of fish in the market. The inference of this study is essential to all relevant stakeholders in the aquaculture industry. The use of Maggot meal if utilized will be a great relief to fish farmers in Nigeria and most developing countries in sub-Saharan Africa that depend on expensive imported fish feed. Maggot meal has proven to provide optimum growth, development and nutrient utilization of *C. gariepinus*. However, more research is needed in the commercial production of maggot meal.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Sogbesan OA, Ugwumba AAA. Bionomics evaluation of garden snail (*Limicolaria aurora*, Jay, 1973; Gastropoda: Limacalaria) meat meal in diet of *Clarias gariepinus* fingerlings (Burchell, 1822); 2006.
2. Ogunji JO, Rahat-UI-Ain ST, Schulz C, Kloas W. Growth performance, nutrient utilization of Nile tilapia, *Oreochromis niloticus* fed housefly maggot meal (magmeal) diets. Turk. J. Fish. Aquat. Sci. 2008a;8:141–147.
3. Ajani EK, LC. Nwanna BO. Musa. Replacement of fishmeal with maggot meal in the diet of Nile Tilapia, *Oreochromis niloticus*, ' World Aquaculture. 2004;35(1): 52-54.
4. FAO. The Effect of COVID-19 on Fisheries and Aquaculture in Asia; 2020. Available: <https://reliefweb.int/sites/reliefweb.int/files/resources/CA9545EN.pdf> (accessed on 21 October 2020).
5. Sogbesan AO, Ajounu ND, Madu CT, Omojowo FS, Ugwumba AAA. Growth response, feed conversion rate and cost benefit of hybrid catfish fed maggot meal-based diet in outdoor tank, 2004 FISON Conference Proceeding. 2005; 816.
6. Okore OO, Ekedo CM, Ubiaru PC, Uzodinma K. Growth and haematological studies of African catfish (*Clarias gariepinus*) juveniles fed with Housefly larva (*Musca domestica*) as feed supplement. International Journal of Agriculture and Earth Science. 2016;2(3): 21-30.
7. Arong GA, Eyo VO. Evaluation of housefly (*Musca domestica*) maggot meal and termite (*Macrotermes subhyalinus*) meal as supplementary feed for African catfish *Clarias gariepinus* (Burchell, 1822). International Journal of Entomology and Nematology. 2017;3(1):42-50.
8. AOAC. Official methods of analysis. 15th edition Association of Official Analytical Chemists, Arlington, VA. 1990;1298.
9. Brett JR, Groves TD. Physiological energetic. In: Fish physiology; 1979.

10. Idowu AB, Amusan AAS, Oyediran AG. The response of *Clarias gariepinus* Burchell 1882) to the diet containing housefly maggot, (*Musca domestica*. Nigerian Journal of Animal Production. 2003;30(1):139- 14.
11. Ogunji JO, Kloas W, Wirth M, Neumann N, Pietsch C. Effect of housefly maggot meal (magmeal) diets on the concentration plasma gluucose, cortisol and blood characteristics of *Oreochromis niloticus* fingerlings. J. Anim. Physiol. Anim. Nutr. 2008b;92:511–518.
12. Akinwande AA, Ugwumba AA, Ugwumba OA. Effects of replacement of fish meal with maggot meal in the diet of *Clarias gariepinus* (Burchell, 1822) fingerlings, The Zoologistl. 2005;(2):41- 46.
13. Dada AA, Akinwande AA. The relationships between stocking density and welfare in farmed rainbow trout. Journal of Fish Biology. 2005;61:493-531.
14. Fasakin AB, Falayi AA, Eyo, Inclusion of poultry manure in the diet of Nile tilapia (*Oreochromis niloticus*), Journal of Fish Technology. 2000;2:51-60.
15. Ekeocha AH, Aganga AA, Emerue PC, Makinde AO. The effect of substituting soybean with groundnut cake on the physiochemical characteristics of novogen cockerel humerus and femur. JASAE. 2023;19(02):1434-1443.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://prh.ikpress.org/review-history/12533>