International Journal of Biochemistry Research & Review



Volume 33, Issue 6, Page 596-603, 2024; Article no.IJBCRR.128170 ISSN: 2231-086X, NLM ID: 101654445

# Exploring Indigenous Legumes of Indonesia: Pigeon Pea (*Cajanus cajan*) and Evaluating Its Impacts on Bone Tissue Structure in Female Rats

### Cicilia Novi Primiani <sup>a,b</sup>, Rafaella Chandraseta Megananda <sup>c</sup> and Titin Kuntum Mandalawati <sup>d\*</sup>

 <sup>a</sup> Department of Biological Education, Faculty of Teacher Training and Education, Universitas PGRI Madiun, East Java, Indonesia.
<sup>b</sup> Department of Pharmacy, Faculty of Health and Science, Universitas PGRI Madiun, East Java, Indonesia.
<sup>c</sup> Department of Food and Agriculture Product Technology, Faculty of Agriculture Technology, Gadjah Mada University, Yogyakarta, Indonesia.
<sup>d</sup> Department of Sport Science, Faculty of Health and Science, PGRI Madiun University, Madiun, East Java, Indonesia.

#### Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: https://doi.org/10.9734/ijbcrr/2024/v33i6939

**Open Peer Review History:** 

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: https://www.sdiarticle5.com/review-history/128170

Original Research Article

Received: 15/10/2024 Accepted: 17/12/2024 Published: 24/12/2024

#### ABSTRACT

**Aims:** The aim of the research was to analyze isoflavone levels and the estrogenic potential of pigeon pea (*Cajanus cajan*) on bone tissue structure preclinically. **Study Design:** Analysis of genistein and daidzein pigeon pea powder using the HPLC method.

\*Corresponding author: E-mail: titin@unipma.ac.id;

**Cite as:** Primiani, Cicilia Novi, Rafaella Chandraseta Megananda, and Titin Kuntum Mandalawati. 2024. "Exploring Indigenous Legumes of Indonesia: Pigeon Pea (Cajanus Cajan) and Evaluating Its Impacts on Bone Tissue Structure in Female Rats". International Journal of Biochemistry Research & Review 33 (6):596-603. https://doi.org/10.9734/ijbcrr/2024/v33i6939.

Preclinical tests were carried out using rat. 24 female white Sprague Dawley rats aged 8-9 months body weight at the start of treatment 150-180 grams. Rats were divided into 3 treatment groups. Control (P<sub>0</sub>), treatment given a solution of pigeon pea seed powder 24 grams: 24 ml of distilled water (P<sub>1</sub>), and treatment given a solution of pigeon pea seed powder 8 grams: 24 ml of distilled water. The solution administration treatment was for 36 days. On the 37th day, surgery was performed and blood and bones were taken. Blood 17  $\beta$  were analyzed using the HPLC method. HE staining bone micro technique.

**Result:** Genistein and daidzein pigeon pea were 247.89828  $\mu$ g/g and 188.61309  $\mu$ g/g respectively. 17 $\beta$ -estradiol P<sub>0</sub> level 237.231 pg/ml; P<sub>1</sub> 603.926 pg/ml P<sub>2</sub> 486.153 pg/ml. Increased proliferation of osteoblasts and fibroblasts in bone tissue structures.

**Conclusion:** Genistein and daidzein are isoflavone compounds found in pigeon pea seeds. Genistein and daidzein can increase  $17\beta$ -estradiol levels. Pigeon pea seeds have the potential to act as natural estrogen.

Keywords: Pigeon pea; osteoblast; daidzein; genistein.

#### 1. INTRODUCTION

Osteoporosis in women is a factor caused by decreased estrogen levels, this results in a decrease in bone mineral density (BMD) [1,2]. Estrogen deficiency causes rapid bone loss within 2-3 years post menopause [3]. The use of hormone replacement therapy (HRT) is a choice for many menopausal women to overcome this [3,4-6].

Continuous use of hormone replacement therapy (HRT) for a long time can pose health risks, making the condition of osteoporosis increasingly complex. Therefore, there is a need for preventive measures and good practices in dealing with osteoporosis. Efforts to prevent osteoporosis can be made by administering natural ingredients containing estrogen [7-9]. The use of natural ingredients is a safer and more effective preventative measure than treatment using synthetic ingredients [9,10,11].

The diversity of local Leguminoceae plants in Indonesia is a wealth of biological natural resources that have not been utilized optimally for health. Leguminoceae are plants that are grouped as phytoestrogens, because they contain high levels of isoflavone components (Yoo et al., 2005), [12-14]. Isoflavones as a compound with a chemical structure similar to 17  $\beta$  estradiol can bind to estrogen receptors [15-17]. The largest components of isoflavones are daidzein, genistein, glycitein, and coumestrol which are thought to be found in Leguminoceae which are distributed in plants and their products [12,14,18].

The chemical structure of isoflavones is similar to 17  $\beta$  estradiol, so its physiological properties resemble the hormone estrogen. Based on its physiological properties, isoflavones have been

widely developed in the food and health sectors. Isoflavone can be used for prevention and treatment, as Hormone Replacement Therapy (HRT) [19,20]. Isoflavones can be used for the treatment of bone fragility, prevention of osteoporosis in postmenopausal women, therapy and prevention of reduction in bone density in postmenopausal women [21-24].

Pigeon pea (*Cajanus cajan*), one of Indonesia's local Leguminoceae, grows abundantly in the lowlands, at an altitude of around 1,800 – 2,000 m. Pigeon pea nutrition 22.6 g/100 g, with glutamic acid, aspartic acid and lysine [25]. The abundance of pigeon peas has not been explored much, the health benefits of pigeon peas have not been explored.

Therefore, pigeon pea is suspected to contain isoflavone compounds, it is necessary to carry out an in-depth analysis, so that pigeon peas, a local bean in Indonesia, can be developed as a natural estrogenic ingredient, especially for preventing osteoporosis. Based on these problems, an analysis is needed to promote natural alternatives side by side reduced dependence on synthetic hormones at affordable solutions. The aim of this study was to analyze isoflavone levels and the estrogenic potential of pigeon pea (*Cajanus cajan*) in bone tissue structure preclinically. It is hoped that the research results will show that pigeon pea can be developed to prevent osteoporosis.

#### 2. MATERIALS AND METHODS

Dried black pigeon pea seeds were obtained from pigeon pea plantations in Ponorogo Regency, East Java, Indonesia. Identification based on Number 0269/Taxo-Plant/Biology/ XI/2023.

#### 2.1 Making Pigeon Pea Seed Powder

500 grams of pigeon pea seeds were dried in an incubator at 60°C for 3 days, then ground using a blender, then ground using a 40 mesh sieve. The fine dry powder of pigeon pea was weighed by total weight.

#### 2.2 Making Pigeon Pea Seed Extract

100 grams of pigeon pea powder was dissolved in 96% ethanol, the solution was homogenized with a vortex for 10 minutes. The solution was left for 30 minutes. Then filtered using Whatman filter paper no. 42, the filtrate was taken. The filtrate that was obtained was then centrifuged at a speed of 8000 rpm for 20 minutes, then 1 ml of the supernatant was taken and then dissolved in 10 ml of distilled water.

#### 2.3 Analysis of Genistein and Daidzein from Pigeon Pea Seed Extract using HPLC Method

Pigeon pea seed powder extract was purified through solid phase extraction using HPLC Shimadzu C18 Sep-Pak, system controller: SCL 10 AVP; solvent delivering unit LC 20 AT; column oven CTO 10 ASVP; SPD 20A UV-Vis detector; column temperature: 25°C; mobile phase: acetonitrile 20% in acetic acid 3%; mobile phase method: isocratic; flow rate: 0.8 ml/min; injection volume: 20 µl detector wavelength: 261 nm; run time: 60 minutes.

#### 2.4 Serum 17β-Estradiol Analysis HPLC Method

On the 37th day, blood was drawn. The  $17\beta$ estradiol analysis procedure was according to Gatti et al. [26] and Yamada et al. [27].

#### 2.5 Grouping of Experimental Animals

Rats were divided into three treatment groups. The first group was a control ( $P_0$ ), the second group ( $P_1$ ) was given a solution of pigeon pea seed powder in the ratio of seed powder: distilled water (24 grams of powder:24 ml of distilled water) and the third group ( $P_2$ ) was given a solution of pigeon pea seed powder in the ratio of seed powder:distilled water (8 grams of powder:24 ml of distilled water).

#### 2.6 Treatment of Experimental Animals

24 female white Sprague Dawley rats aged 10-12 months (premenopause phase) were obtained from the Animal Breeding Unit Blitar, East Java, body weight at the start of treatment 150-180 grams, kept in group cages in the Biosciences experimental animal laboratory at Brawijaya University. Rats were kept in group cages at room temperature (± 27° C), humidity 50-60%, 12 hour light cycle, daily feeding before treatment at 07.00 am, milk pellet composition A with a composition consisting of 12% water, 16% crude protein, crude fat 3-7%, crude fiber 8%, ash 10%, calcium 0.9%-1.2%, phosphorus 0.6%-1% with raw materials yellow corn, wheat bran, SBM. molasses, palm olein, essential amino acids, essential minerals, premixes, and vitamins. Providing water ad libitum. Husk replacement and cage maintenance are carried out every day. Rats were treated by direct induction into the stomach using a gavage tube with pigeon pea solution once a day for 36 days. On the 37th day of the rat, blood was drawn, surgery and bones were taken.

#### 2.7 Making Histological Preparations for HE Staining

Includes the stages of fixation, washing, dehydration, clearing, infiltration, blocking, trimming, sectioning, mounting and staining. Procedure for making histological preparations [28,29]. Changes in bone tissue structure were observed using an optilab microscope.

#### 2.8 Data Analysis

The data obtained, namely data on genistein and daidzein levels from the HPLC test, were analyzed based on the chromatogram results, sample curve area and retention time. Data on bone tissue structure preparations were analyzed for changes in bone tissue structure, osteoblasts and osteoclasts.

#### 3. RESULTS AND DISCUSSION

The results of High Performance Liquid Chromatography (HPLC) analysis of isoflavones (genistein and daidzein) showed genistein and daidzein levels of 247.89828  $\mu$ g/g and 188.61309  $\mu$ g/g, respectively. The results of the HPLC analysis are in Fig. 1.

HPLC analysis Fig. 1 and Table 1, based on sample weight, retention time, and sample curve area.

Isoflavones, lignans, stilbens, koumestans have an aromatic ring structure similar to 17

 $\beta$ -estradiol [30-33], this is often referred to as estrogen like molecules [34,35]. Therefore, these compounds are often called phytoestrogens. Genistein and daidzein are the main isoflavones, which are aglycones that are widely distributed in plants of the Leguminoceae family [33].

Genistein and daidzein are thought to be able to bind to estrogen receptors on ER  $\alpha$  and ER  $\beta$  in the body system [36,37]. The phytoestrogen compounds genistein and daidzein have a mechanism similar to estradiol so they can provide potency in organ systems. Fig. 3 shows that giving pigeon pea powder solution to mice for 36 days was able to increase the proliferation of osteoblasts and fibroblasts in bone tissue structures.

One way ANOVA statistical analysis shows that the significance value (Sig.) 0.004 is smaller than the  $\alpha$  value 0.01, this means that Ho is rejected (Table 2). The results of this statistical analysis stated that given a solution of pigeon pea seed powder had a very significant effect on the number of osteoblasts.



Fig. 1. HPLC with specifications: Shimadzu, injection volume 20 μl, running time 60 minutes, wavelength detector 261 nm, column temperature 25°C, column C 185 μm Shimadzu 120x4.6 mm, mobile phase acetonitrile 20% in acetic acid 3%, Flow rate: 0.8 ml/min. HPLC pigeon pea contained genistein 247,898 μg/g and daidzein 188,613 μg/g

Sample name	Sample weight (g)	RT (min)	Sample curve area	Result (µg/g)	Compound
Pigeon pea	5,0003	37,875	83,28266	188,61309	Daidzein
	5,0003	52,394	74,22387	247,89828	Genistein









Fig. 3. Bone tissue structure of white rat, HE staining, 100x P<sub>0</sub> The structure of bone tissue in normal treatment; P<sub>1</sub> Osteoblasts experience proliferation and increase in fibroblast tissue; P<sub>2</sub> Osteoblasts experience proliferation although not all of them proliferate, there are some osteoclasts

Table 2. One way ANOVA analysis of given solution of pigeon pea seed powder to osteoblasts

	Sum of Squares	Df	Mean Square	F	Sig.	
Between Groups	452.177	2	220.094	7.273	.004	
Within Groups	641.458	21	30.261			
Total	1093.635	23				

Osteoblasts are part of the mesenchymal cells that are responsible for bone formation and development. Osteoblasts produce osteoprotegerin (OPG) as a receptor to prevent bone loss, while 17B-estradiol can stimulate OPG production [38,39]. Genistein and daidzein pigeon pea solution can increase the binding of estrogen receptors so that they can increase bone mineral density (BMD) [40,41]. The hormone estrogen can inhibit bone resorption, suppress the production of IL-1. IL-6 and  $TNF\alpha$ . and inhibit RANK-RANKL interactions by stimulating stromal cells to produce OPG [42,43].

Genistein and daidzein in pigeon pea are phytoestrogens, and can act as substitutes for the hormone estrogen. Therefore, these two phytoestrogen compounds can act as estrogen hormones, especially as estrogen replacement therapy in menopausal conditions to prevent osteoporosis.

#### 4. CONCLUSION

Genistein and daidzein pigeon pea were 247.89828  $\mu$ g/g and 188.61309  $\mu$ g/g respectively. 17 $\beta$ -estradiol P<sub>0</sub> level 237.231 pg/ml; P<sub>1</sub> 603.926 pg/ml P<sub>2</sub> 486.153 pg/ml. Giving pigeon pea can increase 17 $\beta$ -estradiol levels in mice, this can cause proliferation of osteoblasts and fibroblasts of bone tissue

structures. Pigeon pea has the potential to act as a natural estrogen.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

#### ACKNOWLEDGEMENTS

The authors would like to thank the Ministry of Research, Technology, and Higher Education of Indonesia for the financial support of this research. Finally, our great appreciation goes to Muhammad Arisandi for helping in data analysis.

#### **COMPETING INTERESTS**

Authors have declared that no competing interests exist.

#### REFERENCES

1. Levin VA, Jiang X, Kagan R. Estrogen therapy for osteoporosis in the modern era. Osteoporosis International. 2018;29:1049-1055.

- Gosset A, Pouillès JM, Trémollieres F. Menopausal hormone therapy for the management of osteoporosis. Best Practice & Research Clinical Endocrinology & Metabolism. 2021;35(6): 101551.
- Gosset A, Pouillès JM, Trémollieres F. Menopausal hormone therapy for the management of osteoporosis. Best Practice & Research Clinical Endocrinology & Metabolism. 2021;35(6): 101551.
- Gambacciani M, Levancini M. Hormone replacement therapy and the prevention of postmenopausal osteoporosis. Menopause Review/Przegląd Menopauzalny. 2014;13 (4):213-220.
- Eastell R, Rosen CJ, Black DM, Cheung 5. MH. AM. Murad Shoback D Pharmacological management of osteoporosis in postmenopausal women: an Endocrine Society clinical practice Journal auideline. The of Clinical Endocrinology & Metabolism. 2019;104(5): 1595-1622.
- Rozenberg S, Al-Daghri N, Aubertin-Leheudre M, Brandi ML, Cano A, Collins P, Harvey NC. Is there a role for menopausal hormone therapy in the management of postmenopausal osteoporosis?. Osteoporosis international. 2020;31:2271-2286.
- 7. Qin L, Zhang G, Shi Y, Lee K, Leung P. Prevention and treatment of osteoporosis with traditional herbal medicine. In Current topics in osteoporosis. 2005;513-531.
- Martiniakova M, Babikova M, Omelka R. Pharmacological agents and natural compounds: available treatments for osteoporosis. Journal of Physiology & Pharmacology. 2020;71(3).
- 9. Karimi SM, Bayat M, Rahimi R. Plantderived natural medicines for the management of osteoporosis: A comprehensive review of clinical trials. Journal of Traditional and Complementary Medicine. 2024;14(1):1-18.
- 10. Pohl F, Kong Thoo Lin P. The potential use of plant natural products and plant extracts with antioxidant properties for the prevention/treatment of neurodegenerative diseases: in vitro, in vivo and clinical trials. Molecules. 2018;23(12):3283.
- 11. Scott SE, Rozin P, Small DA. Consumers prefer "natural" more for preventatives than for curatives. Journal of Consumer Research. 2020;47(3):454-471.

- Nikolić IL, Savić GIM, Tačić AD, Savić IM. Classification and biological activity of phytoestrogens: A review. Advanced technologies. 2017;6(2):96-106.
- 13. Das S, Sharangi AB, Egbuna C. Jeevanandam J, Ezzat SM, Adetunji CO, Onyeike PC. Health benefits of isoflavones found exclusively of plants of the fabaceae family. Functional Foods and Nutraceuticals: Bioactive Components, Formulations and Innovations. 2020;473-508.
- 14. Whitten PL, Kudo S, Okubo KK. Isoflavonoids. In Handbook of plant and fungal toxicants. 2020;117-137.
- 15. Vitale DC, Piazza C, Melilli B, Drago F, Salomone S. Isoflavones: estrogenic activity, biological effect and bioavailability. European journal of drug metabolism and pharmacokinetics. 2013;38:15-25.
- Křížová L, Dadáková K, Kašparovská J, Kašparovský T. Isoflavones. Molecules. 2019;24(6): 1076.
- 17. Fuentes N, Silveyra P. Estrogen receptor signaling mechanisms. Advances in protein chemistry and structural biology. 2019;116:135-170.
- Sun W, Shahrajabian MH, Cheng Q. Bioactive ingredients of legumes. In Functional foods and nutraceuticals in metabolic and non-communicable diseases. 2022;371-382.
- 19. Larkin T, Price WE, Astheimer L. The key importance of soy isoflavone bioavailability to understanding health benefits. Critical reviews in food science and nutrition. 2008;48(6):538-552.
- Pabich M, Materska M. Biological effect of soy isoflavones in the prevention of civilization diseases. Nutrients. 2019;11 (7):1660.
- Perna S, Peroni G, Miccono A, Riva A, Morazzoni P, Allegrini P, Preda S, Baldiraghi V, Guido D, Rondanelli M. Multidimensional Effects of Soy Isoflavone by Food or Supplements in Menopause Women: A Systematic Review and Bibliometric Analysis. Nat. Prod. Commun. 2016;11:1733–174.
- 22. Lambert MNT, Hu LM, Jeppesen PB. A systematic review and meta-analysis of the effects of isoflavone formulations against estrogen-deficient bone resorption in periand postmenopausal women. Am. J.Clin. Nutr. 2017;106:801–811.
- 23. Akhlaghi M, Ghasemi Nasab M, Riasatian M, Sadeghi F. Soy isoflavones prevent

bone resorption and loss, a systematic review and meta-analysis of randomized controlled trials. Crit. Rev. Food Sci. Nutr. 2020:60:2327–2341.

- 24. Sansai K, Na Takuathung M, Khatsri R, Teekachunhatean S, Hanprasertpong N, Koonrungsesomboon N. Effects of isoflavone interventions on bone mineral density in postmenopausal women: A systematic review and meta-analysis of randomized controlled trials. Osteoporos. Int. 2020;31:1853–1864.
- Putra ID, Marsono Y, Indrati R. Effect of simulated gastrointestinal digestion of bioactive peptide from pigeon pea (Cajanus cajan) tempe on angiotensin-I converting enzyme inhibitory activity. Nutrition & Food Science. 2021;51(2):244-254.
- Gatti R, Gotti R, Gioia MG, Cavrini V. HPLC analysis of pharmaceutical estrogens in raw materials and dosage forms. Journal of pharmaceutical and biomedical analysis. 1998; 17(2):337-347.
- Yamada H, Yoshizawa K, Hayase T. Sensitive determination method of estradiol in plasma using high-performance liquid chromatography with electrochemical detection. Journal of Chromatography B. 2002;775(2):209-213.
- Feldman AT, Wolfe D. Tissue processing and hematoxylin and eosin staining. Histopathology: methods and protocols. 2014;31-43.
- 29. Sampedro-Carrillo EA. Sample preparation and fixation for histology and pathology. Immunohistochemistry and Immunocytochemistry: Methods and Protocols. 2022;33-45.
- 30. Dixon RA. Phytoestrogens. Annu. Rev. Plant Biol. 2004;55(1):225-261.
- Ahmad Hairi H, Jamal JA, Aladdin NA, Husain K, Mohd Sofi NS, Mohamed N, Shuid AN. Demethylbelamcandaquinone B (Dmcq B) is the active compound of Marantodes pumilum var. alata (Blume) Kuntze with osteoanabolic activities. Molecules. 2018;23(7):1686.
- 32. Lee TY, Tseng YH. The potential of phytochemicals in oral cancer prevention and therapy: a review of the evidence. Biomolecules. 2020;10(8):1150.
- Chavda VP, Chaudhari AZ, Balar PC, Gholap A, Vora LK. Phytoestrogens: Chemistry, potential health benefits, and

their medicinal importance. Phytotherapy Research; 2024.

- 34. Vaya J, Tamir S. The relation between the chemical structure of flavonoids and their estrogen-like activities. Current Medicinal Chemistry. 2004;11(10):1333-1343.
- Kiyama R. Estrogenic flavonoids and their molecular mechanisms of action. The Journal of Nutritional Biochemistry. 2023;114:109250.
- Dhananjaya K, Sibi G, Mallesha H, Ravikumar KR, Awasthi S. Insilico studies of daidzein and genistein with human estrogen receptor α. Asian Pacific Journal of Tropical Biomedicine. 2012;2(3):S1747-S1753.
- 37. Chan KK, Siu MK, Jiang YX, Wang JJ, Leung TH. Ngan HY. Estrogen receptor modulators genistein, daidzein and ERB-041 inhibit cell migration, invasion, proliferation and sphere formation via modulation of FAK and PI3K/AKT signaling in ovarian cancer. Cancer cell international. 2018;18:1-14.
- 38. Rickard DJ, Monroe DG, Ruesink TJ, Khosla S, Riggs BL, Spelsberg TC. Phytoestrogen genistein acts as an estrogen agonist on human osteoblastic cells through estrogen receptors  $\alpha$  and  $\beta$ . Journal of cellular biochemistry. 2003;89 (3):633-646.
- An J, Yang H, Zhang Q, Liu C, Zhao J, Zhang L, Chen B. Natural products for treatment of osteoporosis: The effects and mechanisms on promoting osteoblastmediated bone formation. Life sciences. 2016;147:46-58.
- 40. Abdi F, Alimoradi Z, Haqi P, Mahdizad F. Effects of phytoestrogens on bone mineral density during the menopause transition: a systematic review of randomized, controlled trials. Climacteric. 2016;19(6): 535-545.
- 41. Nayeem F, Chen NW, Nagamani M, Anderson KE, Lu LJW. Daidzein and genistein have differential effects in decreasing whole body bone mineral density but had no effect on hip and spine density in premenopausal women: A 2year randomized, double-blind, placebocontrolled study. Nutrition Research. 2019;68:70-81.
- 42. Drugarin D, Drugarin M, Negru S, Cioace R. RANKL/RANK/OPG molecular complexcontrol factors in bone remodeling. TMJ. 2003;53(3-4):297-302.

## 43. Hooshiar SH, Tobeiha M, Jafarnejad S. Soy isoflavones and bone health: Focus on

the RANKL/RANK/OPG pathway. BioMed research international. 2022;(1):8862278.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history: The peer review history for this paper can be accessed here: https://www.sdiarticle5.com/review-history/128170