



# Knowledge, Perception Regarding the Effect of Tobacco Use and Cessation Practices: A Case Study from College of Health Technology Students, Lagos State, Nigeria

Oyekan Mojisola Adejoke <sup>a\*</sup>, Whenayon Sanusi <sup>b</sup>,  
Omotoso Ayodele Jacob <sup>c</sup>, Raji Kazeem Ajayi <sup>c</sup>  
and Odeleye Jonathan Oluwafemi <sup>c</sup>

<sup>a</sup> Lagos State College of Health Technology, Nigeria.

<sup>b</sup> Lagos State Health District I (Ikotun Elf PHC), Nigeria.

<sup>c</sup> Edo State College of Health Sciences and Technology, Nigeria.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

DOI: <https://doi.org/10.9734/ajarr/2025/v19i1866>

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

<https://www.sdiarticle5.com/review-history/127866>

**Original Research Article**

**Received: 20/10/2024**

**Accepted: 23/12/2024**

**Published: 09/01/2025**

## **ABSTRACT**

The study is the investigation of knowledge, perception regarding the effect of tobacco use and cessation practices among students of Lagos state college of health technology in Yaba mainland local government area. Snowball sampling techniques selected by Lagos State, Nigeria, and 200

\*Corresponding author: Email: [mojisola.adejokeoyekan@gmail.com](mailto:mojisola.adejokeoyekan@gmail.com);

**Cite as:** Adejoke, Oyekan Mojisola, Whenayon Sanusi, Omotoso Ayodele Jacob, Raji Kazeem Ajayi, and Odeleye Jonathan Oluwafemi. 2025. "Knowledge, Perception Regarding the Effect of Tobacco Use and Cessation Practices: A Case Study from College of Health Technology Students, Lagos State, Nigeria". *Asian Journal of Advanced Research and Reports* 19 (1):103-119. <https://doi.org/10.9734/ajarr/2025/v19i1866>.

respondents. A structured and validated questionnaire was used to elicit information from the respondents.

The study adopted a descriptive cross-sectional study design. Demographic data were analyzed using frequencies and percentages in tabular form to test demographic data of the respondents with chi-square ( $\chi^2$ ) analysis was used to test the hypothesis at 0.005 level of significance.

The result revealed that majority of the students knows that tobacco use has harmful effects on its users, while few have a perception about tobacco use cessation techniques and majority have the positive perception that role modeling of health professionals on cessation practices plays an important role in patients quitting tobacco use.

Based on these findings, some recommendations were suggested among which are the urgent need of Government to enhance the synergy between the Ministry of Health and Ministry of Education so as to include cessation techniques in the training curricula of health professionals as well as the development of improved and effective enforcement and compliance on cigarette ban in public places.

*Keywords: Knowledge; perception; tobacco use and cessation practices.*

## 1. INTRODUCTION

Tobacco use is one of the major preventable causes of untimely death, disease, and disability, which accounts for a significant percentage of various health inequalities around the earth. (Ezzati, M al 2002). It is ranked 6 out of 8 top causes of death in the world. Tobacco use comprises of the smoking and non-smoking tobacco products; Tobacco smoking is a custom harmful to the eyes, odious to the nose, dangerous to the brain and harmful to the lungs. It poses colossal health and non-health related costs to the affected persons and society at large.

In many modern societies, smoking tobacco is an accepted habit with adults of all ages and sexes. The practice used to be socially acceptable in Europe and the United States but is less so now (Brar et al., 2022). Laws have been passed in some American States restricting smoking, and people can no longer smoke in public places like bars, restaurants, and clubs. (Shafey et al, 2012)

In Africa, similar laws that are supposed to be in effect have remained on paper, and there is minimal enforcement where people can or cannot smoke (Weiss et al., 2020; Gallaway et al., 2019).

Therefore, a lower awareness of the dangerous effects of smoking is just one of the many reasons for the higher predominance of smoking among lower socio-economic groups. A successful strategy for controlling tobacco use for disadvantaged groups should address their living conditions and social environment and inform them about the effects of smoking on health.

## 2. METHODOLOGY

### 2.1 Research Design

A descriptive research design was adopted in conducting this study. It was used to carry out design in order to allow relevant instruments to be identified, evaluated and summarized.

### 2.2 Study Design

The study was a descriptive cross-sectional study design to assess the knowledge, perception regarding the effect of tobacco use and cessation practices among students of Lagos State College of Health Technology in Yaba local government area.

### 2.3 Study Population

The study population comprises of individuals who use tobacco and are students of Lagos State College of Health Technology.

### 2.4 Sample Size Determination

The sample size (n) was calculated using the Cochran's sample size formula

$$n = \frac{Z^2 pq}{d^2} \quad (\text{Cochran WG, 1997})$$

Where n = minimum sample size when target population  $n \geq 10,000$

z= standard normal deviate at 95% confidence = 1.96

p=the proportion in the target population estimated to be current tobacco users in Nigeria = 16.7% (GATS, 2012)

q= 1-p=proportion of the target population estimated to be non-tobacco users: 1-0.167=0.833  
 d=margin of error=5%=0.05

Applying this formula:

$$n = \frac{z^2 pq}{d^2}$$

$$n = \frac{z^2 pq}{d^2} = \frac{1.96^2 \times 0.167 \times 0.833}{(0.05)^2}$$

$$= \frac{3.84 \times 0.167 \times 0.833}{0.0025}$$

$$= \frac{0.534}{0.0025}$$

$$n = 213 \text{ respondents (smokers)}$$

With reference to population under investigation

N = Population of students in the College of Health Technology = 1,193  
 n= Calculated minimum sample size from the Cochran formula  
 n<sub>f</sub> = Calculated sample size relative to study population

$$n_f = \frac{n}{\text{Expected response rate}}$$

$$n_f = \frac{n}{1 + \frac{n}{N}} \quad (1)$$

$$= \frac{213}{1 + \frac{213}{1193}}$$

$$= \frac{213}{1 + 0.18}$$

$$= \frac{213}{1.18}$$

$$= 180 \text{ respondents (smokers)}$$

Adjusting for non- response

To compensate for questionnaire that may not be completely filled will be set at 10%

$$na = \frac{n_f}{\text{Anticipated response rate}}$$

$$na = \frac{n_f}{0.90}$$

$$= \frac{180}{0.90}$$

$$= 200 \text{ respondents (smokers)}$$

## 2.5 Sampling Methodology

### 2.5.1 Methodology

The study is a multi-stage sampling method and a sample frame designed for each level of students, and the respondents were selected using the snowball sampling method: The snowball sampling method is a non-probability sampling technique used to identify potential respondents (i.e., smokers); whereby participants identified and invited other possible potential respondents.

### 2.5.2 Data collection tool and techniques

A pre-tested, structured interviewer-administered questionnaire developed from modified version of the global health professional surveys core questionnaire which was developed from global adult tobacco survey and validated by collaboration of centres of Disease Control and Prevention (CDC), World Health Organization, CDC Foundations, John Hopkins Bloomberg School of Public Health (JHSPH) and Research Triangle Institute (RTI) will be used for the data collection. The questionnaire was designed to obtain information.

### 2.5.3 Data collection

The data was collected among students of Lagos State College of Health Technology in July 2016 after modification and adjustment of the pre-tested questionnaire as a result of discrepancies obtained during the pre-test. The researcher collected data and assisted by the 8 participants (respondents).

### 2.5.4 Data analysis

The data collected were analyzed using the Statistical Package for Social Sciences version 20.

### 2.5.5 Data presentation

Quantitative data are presented with frequency tables and charts, percentages, means, and standard deviation were determined, chi-square was used to compare variables confidence, the interval the was set at 95%, and statistical tests were considered significant at p values less than 0.05.

## 2.6 Ethical Consideration

The purpose and nature of the study were communicated to the respondents, written

informed consent was obtained from the students before the participation in the study, and anonymity of respondents was maintained by non-use of identifiers and personal information. They were also informed that they were free not to participate and or opt out at any time.

### 3. RESULTS AND DISCUSSION

#### 3.1 Results and Discussion of Findings

The overall percentage grading of respondents knowledge regarding the health effects of tobacco use was found to be 55.5% which is scored fair knowledge, while 28.7% was scored poor knowledge and I do not know 15.8% neutral on lung, heart, lung and other diseases, low birth weight from pregnant smoker and low sperm count for male sexual dysfunction.

For stroke, HIV, road traffic accidents, 6.3% out of the respondents affirmed that these could be

the health effects of tobacco use and this was rated poor knowledge regarding the health effect of tobacco use, while 67.6% affirmed negative and 59% were neutral; this is rated poor overall knowledge.

As shown in Table 3, 183 subjects representing 91.5% agreed that knowledge regarding the effect of tobacco use would not significantly promote the knowledge, perception regarding the harmful effect of tobacco use and cessation practises among students of LASCOHET in Yaba Mainland Local Government area of Lagos State.

Therefore, it can be concluded that knowledge regarding the effect of tobacco use will not significantly promote the knowledge, perception regarding the harmful effect of tobacco use and cessation practices among students of LASCOHET in Yaba mainland local government area of Lagos state.

**Table 1. Respondents patterns of tobacco use**

Age Range(Yrs) Frequency	Below 14 Daily	14-18 Weekly	19-25 Monthly	26-30 Occasionally	Total
Age of First Smoked	0	178	22	0	200
Percentage (%)	0.0	89.0	11.0	0.0	100%
Responses	195	0.0	0.0	5.0	200
Observed	97.5	0.0	0.0	2.5	100%
Expected	195	0	0	5	200
Percentage(%)	97.5	0.0	0.0	2.5	100%
Average Per Day/Sticks	1	2	3	4-10	Total
Observed	32	7	64	97	200
Percentage %	16	3.5	32	48.5	100%
<b>Average Per Week/Sticks</b>	<b>8-14</b>	<b>15-21</b>	<b>22-28</b>	<b>29-30</b>	<b>Total</b>
<b>Observed</b>	<b>32</b>	<b>7</b>	<b>64</b>	<b>97</b>	<b>200</b>
<b>Percentage%</b>	<b>16</b>	<b>3.5</b>	<b>32</b>	<b>48.5</b>	<b>100%</b>

Mean Age ± S.D 16.7 ± 1.88

**Table 2. Respondents knowledge regarding the effect of tobacco use**

Statement Frequency (100%)	Yes	No	I do not know	Total /(%)
Harmful to health	183(91.5)	17(8.5)	0 (0.0)	200(100)
Second-hand smoke is harmful to health	174(87.0)	10(5.0)	16 (8.0)	200(100)
<b>Diseases associated with smoke:</b>				
Lung Diseases	170(85.0)	13(6.5)	17 (8.5)	200(100)
Heart Diseases	159(79.5)	23(11.5)	18 (9.0)	200(100)
Lung and other cancer	167(83.5)	19(9.5)	14 (7.0)	200(100)
Stroke	26(13.0)	47(23.5)	127(63.3)	200(100)
Sudden Death	69(34.5)	62(31.0)	69(34.5)	200(100)
HIV/AIDS	0(0.0)	176(88.0)	24(12.0)	200(100)
Road Traffic Accident	12 (6.0)	162(81.0)	26(13.0)	200(100)
Low birth weight in female smokers children	15(7.5)	112(56.0)	73(36.5)	200(100)
Low Sperm counts	40(20.0)	73(36.5)	87(43.5)	200(100)

**Table 3. Respondents knowledge regarding the effect of tobacco use**

<b>Responses</b>	<b>yes</b>	<b>no</b>	<b>total</b>
Observed	183	17	200
%	91.5	8.5	100
Expected	100	100	200
%	50	50	100

Agree	183	91.5
Disagree	17	8.5
<b>Total</b>	200	100.0

$\chi^2$  value= 0.000; P 0.05 (significant) df =1

**Table 4. Respondents level of tobacco use**

<b>Age range(yrs)</b>	<b>Below 14</b>	<b>14-18</b>	<b>19-25</b>	<b>26-30</b>	<b>Total</b>
<b>Frequency</b>	<b>Daily</b>	<b>Weekly</b>	<b>Monthly</b>	<b>Occasionally</b>	
Age of first smoked	0	178	22	0	200
Percentage(%)	0.0	89.0	11.0	0.0	100%
Responses	195	0.0	0.0	5.0	200
Observed	97.5	0.0	0.0	2.5	100%
Expected	195	0	0	5	200
Percentage(%)	97.5	0.0	0.0	2.5	100%
Average per day/sticks	1	2	3	4-10	Total
Observed	32	7	64	97	200
Percentage %	16	3.5	32	48.5	100%
<b>Average per week/sticks</b>	<b>8-14</b>	<b>15-21</b>	<b>22-28</b>	<b>29-30</b>	<b>Total</b>
<b>Observed</b>	<b>32</b>	<b>7</b>	<b>64</b>	<b>97</b>	<b>200</b>
<b>Percentage%</b>	<b>16</b>	<b>3.5</b>	<b>32</b>	<b>48.5</b>	<b>100%</b>

Mean Age  $\pm$  S.D 16.7  $\pm$  1.88

**Table 5. Respondents level of tobacco use**

<b>Responses</b>	<b>Yes</b>	<b>know</b>	<b>Total</b>
Observed	183	17	200
%	91.5	8.5	100
Expected	100	100	200
%	50	50	100

	<b>Daily</b>	<b>Occasional</b>	<b>Total</b>	<b>%</b>
Agree	179	4	183	91.5%
Disagree	16	1	17	8.5%
Total	195	5	200	100.0%

$\chi^2$  value= 0.000; P 0.05(significant) df = 3

As shown in Table 5, 183 subjects representing 91.5% agreed that the level of tobacco use cessation practices would not significantly promote the knowledge, perception regarding the harmful effect of tobacco use and cessation practises among students of LASCOHET in Yaba Mainland Local Government area of Lagos State.

On the other hand,17 subjects representing 8.5% disagreed in their opinion on the issue.

The data were subjected to chi-square analysis  $\chi^2$  test for acceptance or non- acceptance of the hypothesis.The obtained chi-square value at df 3, and the significant level of 0.05 is 0.000. Since the obtained P value  $\chi^2$  is less than < 0.05, the hypothesis is rejected. Therefore, it can be concluded that the level of tobacco use cessation practices will significantly promote the knowledge, perception regarding the harmful effect of tobacco use and cessation practises among students of LASCOHET in Yaba Mainland Local Government area of Lagos State.

**Table 6. Perception of tobacco use cessation practices**

Statement	Frequency (100%)		
	Yes	No	IDont Know
Ever tried to stop smoking during the past year,	79 (39.5)	121(60.5)	0 (0.0)
Ever tried to stop using tobacco now	50 (25.0)	71 (35.5)	79 (39.5)
If Yes, Why?	Health reason	It wastes money	To please family
	25 (12.5)	10 (5.0)	15 (7.5)
If No, Why?	I don't want to stop	Nothing bad to stop	Addiction/ Withdrawal
	25 (12.5)	29 (14.5)	17(8.5)
Quit period	Not stopped	<1mnth	1-5mnths
	9 (4.5)	5 (2.5)	4 (2.0)
	1year	2years	> 3years
	2 (2.0)	2 (2.0)	1(0.5)
Help /Advice to stop tobacco use	Program	Friend	Family
	6(3.0)	72 (36.0)	10 (5.0)
	Program/ Professional	Professional/family/friend	None
	6 (3.0)	6 (3.0)	100
<b>Ever heard of Nicotine Replacement Therapy/ others</b>	<b>Yes</b>	<b>No</b>	<b>Don't Know</b>
	20	60	120




Fig. 1. Chart for respondents knowledge regarding effects of tobacco use




Fig. 2. Chart for Table 6 on the perception of tobacco cessation practices among the respondents

**List 1. Grading of respondent’s overall perception of tobacco use cessation practices**

Perception Grade	Frequency	Percentage (%)
Good	79	39.5
Poor	121	60.5
Neutral	0	0.0
<b>Total</b>	<b>200</b>	<b>100.0</b>

**Table 7. Respondents perception of tobacco use cessation practices**

Responses	Yes	no	I do not know	Total
Observed	50	71	79	200
%	25.0	35.5	39.5	100
Expected	66.67	66.67	66.67	200
%	33.33	33.33	33.33	100.0
Agree	50		25.0	
Disagree	71		35.5	
Neutral	79		39.5	
<b>Total</b>	<b>200</b>		<b>100.0</b>	

*X<sup>2</sup> value= 0.035; P 0.05(significant) df = 2*

**Yes denotes good: No denotes poor: I do not know: denotes neutral:** The negative perception regarding respondents on tobacco cessation approaches from health worker was rated poor perception because only 20% (health worker), 10% Prescription drugs, 35% tobacco cessation clinic with good knowledge, hence poor scoring on perception of tobacco use cessation practices; While the positive perception (25%) health workers, 3% prescription drugs, (1%) tobacco cessation clinic with overall percentage scoring of 35,5% positive perception on tobacco use.

**Grading of respondent’s overall perception of tobacco use cessation practices:** 39.5 % of the respondents surveyed had a good perception of tobacco use cessation while 60.5% have a poor perception of tobacco cessation practices.

As shown in Table 7, 50 subjects representing 25%, agreed that perception of tobacco use cessation practices will not significantly promote the knowledge, perception regarding the harmful effect of tobacco use and cessation practices among students of LASCOHET in Yaba Mainland Local Government area of Lagos State.

On the other hand, 71 subjects representing 35.5% disagreed, while 79 subjects are representing 39.5% were neutrals in their opinion on the issue.

The data were subjected to chi-square analysis X<sup>2</sup> test for acceptance or non- acceptance of the hypothesis. The obtained chi-square value at df 2

and the significant level of 0.05 is 0.035. Since the obtained P value X<sup>2</sup> is less than < 0.05, the hypothesis is rejected. Therefore, it can be concluded that perception of tobacco use cessation practices will significantly promote the knowledge, perception regarding the harmful effect of tobacco use and cessation practises among students of LASCOHET in Yaba Mainland Local Government area of Lagos State.

Table 8 Finally, the participants were requested of their readiness to make available assistance to patients who like to stop smoking and 31(15.5%) responded positively in providing help to patients. 138 (69%) responded positively to getting specific training on tobacco cessation techniques and giving advice and information about smoking cessation.

More than three quarter 148 (71%) responded positively to advise patients who use other tobacco products to quit the use. The respondents 127(63.5%) agreed that the chances of patients to quit smoking are high if a health professional advice.

For questions about tobacco education in the health professionals’ curricula, only 33.33% sensed that they are being educated satisfactorily about the health effects of smoking, and tobacco cessation methods. Approximately 50% of the students felt that they should be educated about tobacco cessation methods, including counselling; recent training about tobacco smoking is not methodical and incorporated with other disciplines.

**Table 8. Perception of role modelling in tobacco cessation practices**

Statement	Frequencies (%)		
	Yes	No	I do not know
Should you be involved in the use of any tobacco products	3 (21.5)	67 (33.5)	90 (45.0)
Do you serve as a role model for patients and the public	98 (49.0)	25 (12.5)	77 (38.5)
As a tobacco smoker, is it less likely to advise patients to stop smoking	31 (15.5)	66 (33.0)	103 (51.5)
As a user of other tobacco products (chewing tobacco, snuff, cigars/ pipes), is it likely to advise patients to stop smoking	37 (18.5)	2 (1.0)	1 (0.5)
Should information on tobacco use	147 (23.5)	13 (6.5)	40 (20.0)
Be taken while taking patients' history			
Should you get specific training on Tobacco cessation techniques	138 (69.0)	10(5.0)	52 (26.0)
Should medical professionals be role models by being non-smokers to advise their patients smoking cessation?	99 (50.0)	25 (12.5)	76 (37.5)
Should all medical schools have smoking cessation clinics with facilities for counseling, treatment & follow-up?	146 (22.5)	13 (6.5)	41 (21.0)
Should smoking amongst medical teachers and students be the main obstacle in effectively implementing tobacco education?	40 (20.0)	13 (6.5)	147 (23.5)
Does the current curriculum teach adequately about the health effects of active and passive smoking?	138 (69.0)	10(5.0)	52 (26.0)
Should you routinely advise Patients who smoke to quit	174 (87.0)	26 (13.0)	0.(0.0)
Does the current curriculum teaches about clinical guidelines, tobacco cessation methods and its contraindications.	147 (23.5)	13 (6.5)	40 (20.0)
Should you routinely advice patients who use other tobacco products to quit using these products	148 (74.0)	26 (13.0)	26 (13.0)
Role in giving advice/information on smoking cessation to patients	138 (69.0)	13 (6.5)	39 (19.5)
Are a patient's chances of quitting smoking increased if health Professional advice him or her to quit	127 (63.5)	32 (16.0)	41 (20.5)
Does your religion support the use Of any tobacco products	168 (84.0)	0(0.0)	32 (16.0)
Do you serve as 'role models for patients	195 (97.5)	5 (5.0)	0 (0.0)
Role in advising patients	187(93.5)	9(5)	4(2.0)
Chances of patients quitting	17(85.5)	10(5.0)	19(9.5)

**Table 9. Perception of role modelling in the tobacco cessation practices**

Responses	Yes	Know	I do not know	Total
Observed	99	24	77	200
%	49.5	12.0	38.5	100
Expected	66.67	66.67	66.67	200
%	33.3	33.3	33.3	100

Agree	99	49.5%
Disagree	24	12.0%
Neutral	77	38.5%
<b>Total</b>	<b>200</b>	<b>100.0</b>

$X^2$  value= 0.00; P 0.05(significant) df = 2

**Table 10. Relationship between socio-demographic characteristics and knowledge on the health effect of tobacco use**

Socio-demographic Characteristics	Frequency %		Statistical test
	Good (n=183)	Poor (n=17)	
Age Mean $\pm$ SD	27.2 $\pm$ 4.04		t=29.14 p=0.000
Gender			
Male	157	17	$X^2=110.6$ df =3 p=0.000
Female	26	0	
Course Year			
First Year	15	0	*p=0.000
Second Year	62	0	
Third Year	90	16	
Fourth Year	16	1	
Religion			
Christianity	68	16	*p=0.000
Islam	115	1	
Ethnic Group			
Yoruba	98	1	*p =0.000
Hausa/Fulani	6	0	
Igbo	79	16	
Marital status			
Single	163	15	*p =0.000
Married	20	2	
Separated			
Widowed			

\*Fisher's exact tests

Almost half of the students approved that health professionals curricula should include a distinct module about tobacco education. Three-quarters of the students from agreed about teaching tobacco cessation, and counseling, in a separate module of the health professionals' curriculum.

As shown in Table 9, 99 subjects representing 49.55%, agreed that the perception on role modelling in the tobacco use cessation practices would not significantly promote the knowledge, perception regarding the harmful effect of tobacco use and cessation practises among

students of LASCOHET in Yaba Mainland Local Government area of Lagos State.

On the other hand, 24 subjects representing 12.0% disagreed, while 77 subjects representing 38.5% were neutral in their opinion on the issue.

The data were subjected to chi-square analysis  $X^2$  test for acceptance or non- acceptance of the hypothesis. The obtained chi-square value at df 2, and the significant level of 0.05 is 0.000. Since the obtained P value  $X^2$  is less than  $< 0.05$ , the hypothesis is rejected. Therefore, it can be concluded that the perception on role modelling

in the tobacco use cessation practices will significantly promote the knowledge, perception regarding the harmful effect of tobacco use and cessation practices among students of LASCOHET in Yaba Mainland Local Government area of Lagos State.

Table 10 shows that at 95% confident interval, age has no effect on knowledge regarding health effect of tobacco use, the p-value is 0.000, this is less than 0.05; it means that there is a significant relationship between age and knowledge. It also

shows that there is statistically significant relationship gender and knowledge as 157(78.5%) males know more than the females.

Table 11 at 95% confident interval p-value is 0.000, this is less than 0.05; it means that there is a significant relationship between age and perception of tobacco cessation practices. It also shows that there is a statistically significant relationship between gender, religion, educational background, and the respondents.

**Table 11. Relationship between socio-demographic characteristics and perception of tobacco cessation practices**

Socio-demographic Characteristics	Frequency %		Statistical test
	Good (n=183)	Poor (n=17)	
Age			
Mean $\pm$ SD	27.2 $\pm$ 4.04		t=20.35 p=0.000
Gender			
Male	57	117	X <sup>2</sup> =71.43 df =1 p=0.0003
Female	22	4	
Course Year			
First Year	4	11	*p=0.000
Second Year	38	24	
Third Year	33	73	
Fourth Year	4	13	
Religion			
Christianity	57	74	*p=0.000
Islam	22	47	
Ethic Group			
Yoruba	48	59	*p =0.000
Hausa/Fulani	5	1	
Igbo	37	61	
Marital status			
Single	64	114	*p =0.001
Married	15	7	
Separated			
Widowed			

\*Fisher's Exact tests

**Table 12. Association between pattern and knowledge of the effects of tobacco use Association of a pattern of use and knowledge of the health effects of tobacco use**

Health problems associated	Below 10			Between 11 and 20		
	Y	N	IDK	Y	N	IDK
Tobacco use						
Lung Disease	168	13	12	2	0	5
Heart Diseases	152	23	18	7	0	7
Lung and Other Cancers	163	17	18	4	2	1
Stroke	22	47	124	4	0	3
Sudden Death	69	55	69	0	7	0
HIV/AIDS	0	173	20	0	3	4
Road Traffic Accident	12	155	26	0	7	0
Low Birth Weight	15	108	70	0	4	3
Low Sperm Count	40	66	87	0	7	0

Y = Yes. N= No & IDK = I don't know

**Table 13. Effect of socio-demographic characteristics of respondents on knowledge, perception regarding the effect of tobacco use and cessation practices**

Variable	Test statistics	p values
knowledge on harmful effect of tobacco use	$\chi^2 = 137.78$ df = 1	.000
perception on cessation practices	$\chi^2 = 6.730$ df = 2	.035
perception on role modelling	$\chi^2 = 44.59$ df = 2	.000

Table 10 Majority of the respondent are heavy tobacco users and have good knowledge on the health effect on tobacco use.

Number of cigarettes taken per day below 11

Majority of the respondents are heavy tobacco users but have good knowledge of the health effect of tobacco use.

The P value for knowledge on the health effect of tobacco use was found to be 0.00 and is less than the level of significance 0.05; hence, the respondents have a good knowledge regarding the effect of tobacco use. ( $X^2 = 137.78$ ;  $< .000$ )

It shows that having a positive perception of cessation practices was found to be associated with having good knowledge. The P value was found to be less than the level of significance ( $X^2 = 6.73$ ;  $< 0.35$ ).SS

Positive perception of role modelling was found to be associated with good knowledge. The P value was found to be less than the level of significance. ( $X^2 = 44.59$ ;  $< 0.00$ ).

#### 4. CONCLUSION

The class representations of samples of smokers from the Lagos State College of Health Technology and showed that lower level status is related to less knowledge of the health effects of smoking and toxic elements of smoke. These findings about the consciousness of health effects were consistent with earlier research, awareness of toxins in smoke.

We note the general acceptance of our findings across the other health professional institutions in other countries and hypothesise those alike social variations exist in other high-income countries.

Misconception about nicotine was common with over 40% of all smokers trusting that nicotine

causes most of cancer. Nearly half of the smokers in the college and the majority of smokers with low socioeconomic status fostered this belief. It is significant that such misunderstandings are modified, as smokers who thought that nicotine is carcinogenic are not likely to decide to use nicotine replacement therapies, which are verified treatments for ending.

It is possible that a small number of smokers believed that nicotine played a role in the development of cancer because of its role as an addictive substance that keeps people smoking; further research is required to explore this.

Socioeconomic status of lower-level differences in the knowledge that tobacco smoke has mercury was small because little people were conscious of the presence of this substance in tobacco smoke. Mass media campaigns or cigarette pack cautioning labels have not sufficiently emphasised the fact that tobacco smoke comprises of mercury.

We note that normally education had a stronger influence than income, where knowledge was scarce (for example, knowledge about impotence, cyanide, mercury, and nicotine as a cause of cancer), and a weaker influence than income, where knowledge was widespread (for example, knowledge about heart disease and lung cancer).

The only element of knowledge that did not fit this pattern was carbon monoxide. We speculate that when knowledge about the health effects of a substance is available for the first time, higher education groups gain knowledge faster. They are better able to obtain information and use information technology and have a broader knowledge and access to information sources.

It is prominent that among other health professional institutions, smokers in Canada had the utmost level of consciousness of the harms

of smoking and toxic compounds in smoke. This is likely due to the strong anti-smoking education campaigns in Canada. In 2001, Health Canada announced the commencement of a five-year \$480 million programme to bolster existing tobacco control campaigns.

About 40% of the budget was allocated to mass media campaigns targeting Canadians of all ages. (Canada Tobacco Control, 2004) Canada has one of the most progressive warning labels on cigarette packs in the world. Under Canadian law, 16 rotated picture-based warnings are required in the top 50% of the front and back of the cigarette package. The graphic pictures include a diseased mouth, a lung tumour, and a brain after a stroke, a damaged heart, and a limp cigarette as part of an impotence warning.

Also, there are 16 rotating interior messages. On the side of the package, there is a range of yields for six substances. Cigarette packs in the USA, UK, and Australia contain text warnings. Graphic warnings are expected to be introduced in Australia within two years and in the UK within five years. Package warning labels for health effects and smoke constituents have been linked to greater health knowledge among Canadian, US, UK, and Australian smokers (Hammond et al, 2008).

In addition, the study could investigate the differences in knowledge of the health effects of the components of tobacco smoke; the effect on the health of tobacco smoke in the environment; Myths about smoking (for example exercise or vitamins undo most of the effects of smoking); and the perception of smokers about their risk of developing diseases related to smoking.

#### **DISCLAIMER (ARTIFICIAL INTELLIGENCE)**

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

#### **COMPETING INTERESTS**

Authors have declared that no competing interests exist.

#### **REFERENCES**

Abdullah, A. S., Stillman, F. A., Yang, L., Luo, H., Zhang, Z., & Samet, J. M. (2013). Tobacco

use and smoking cessation practices among physicians in developing countries: A literature review (1987–2010). *International Journal of Environmental Research and Public Health*, 11, 429–455.

Abroms, L. C., Windsor, R., & Simons-Morton, B. (2008). Getting young adults to quit smoking: A formative evaluation of the X-Pack program. *Nicotine & Tobacco Research*, 10(1), 27–33.

Alakija, W. (1984). Smoking habits of medical students of the University of Benin, Nigeria. *Nigerian Medical Journal*, 14, 171–174.

Al-Haqwi, A. I., Tanim, H., & Asery, A. (2016). Knowledge, attitude, and practice of tobacco smoking by medical students in Riyadh, Saudi Arabia.

Almerie, M. Q., Matar, H. E., Salam, M., Morad, A., Abdulaal, M., Koudsi, A., et al. (2008). Cigarettes and water pipe smoking among medical students in Syria: A cross-sectional study. *International Journal of Tuberculosis and Lung Disease*.

Andersson, P., & Leppert, J. (2001). Men of low socioeconomic and educational level possess pronounced deficient knowledge about the risk factors related to coronary heart disease. *Journal of Cardiovascular Risk*, 8, 371–377.

Ashton, C. H., & Kamali, F. (1995). Personality, lifestyles, alcohol, and drug consumption in a sample of British medical students. *Medical Education*, 29, 187–192.

Aslam, S. K., Zaheer, S., & Shafique, K. (2011). Health professional students' perceptions regarding their role in tobacco control: Findings from the Global Health Professional Students Survey, Pakistan. *Substance Abuse Treatment, Prevention, and Policy*, 6(9), 25.

Awopeju, O. F., Erhabor, G. E., Awosusi, B., Awopeju, O. A., & Adewole, O. O. (2010). Smoking prevalence among health professional students of Obafemi Awolowo University, Ile-Ife, and the University of Lagos in southwestern Nigeria. *Annals of Medicine and Health*.

Awotedu, A. A., Jordaan, E. R., Ndukwana, O. Z., Fipazano, F., & Martinez, J. (2006). The smoking habits, attitudes towards smoking, and knowledge regarding anti-smoking legislation of students in an institution of higher learning in Eastern Cape Province of South Africa. *Family Practice*, 48, 14–18.

- Ayanian, J. Z. (1999). Perceived risks of heart disease and cancer among cigarette smokers. *JAMA*, 281, 1019–1021.
- Bande, E. O., & Osadiaye, J. A. (1987). Attitudes and smoking habits of physicians at the Lagos University Teaching Hospital. *Journal of the National Medical Association*, 79, 430–432.
- Bandura, A. (1977). Self-efficacy: Toward a unifying theory of behavioral change. *Psychological Review*, 84, 191–215.
- Bandura, A. (2008). *Social foundations of thought and action: A social cognitive theory*. Englewood Cliffs, NJ: Prentice-Hall.
- Beltre, M., Kebede, D., & Kassaye, M. (2006). Modifiable risk factors for coronary heart disease among young people in Addis Ababa. *East African Medical Journal*, 74(6), 376–381.
- Benson, D., Canchola, J. A., & Catania, J. A. (2000). Random selection in a national telephone survey: A comparison of the Kish, next-birthday, and last-birthday methods. *Journal of Official Statistics*, 16, 53–60.
- Birch, D., Ashton, H., & Kamali, F. (1998). Alcohol drinking, illicit drug use, and stress in junior house officers in the northeast of England. *Lancet*, 352, 785–786.
- Bobak, M., Jha, P., & Nguyen, S. (2000). Poverty and smoking. In P. Jha & F. J. Chaloupka (Eds.), *Tobacco control in developing countries* (pp. xx–xx). Oxford: Oxford University Press.
- Borland, R., & Hill, D. (1997). Initial impact of the Australian tobacco health warnings on knowledge and beliefs. *Tobacco Control*, 6, 317–325.
- Brar, M., Randhawa, A., & Chaudhary, N. (2022). Tobacco use in medical college students and its effect on their attitudes and practices. *Current Practice in Medical Science*, 5(14), 161–170.
- Center for Disease Control and Prevention (CDC). (2011). Vital signs: Current cigarette smoking among adults aged ≥18 years — United States, 2005–2010. *MMWR Morbidity and Mortality Weekly Report*, 60(35), xx–xx.
- Chatkin, J., & Chatkin, G. (2009). Learning about smoking during medical school: Are we still missing opportunities? *International Journal of Tuberculosis and Lung Disease*, 13, 429–437.
- Chatkin, J., & Chatkin, G. (2011). Learning about smoking during medical school: Are we still missing opportunities? *International Journal of Tuberculosis and Lung Disease*, 13, 429–437.
- Christen, A. C., Sawson, B. Z., Glover, E. D., & Henderson, A. H. (1982). Smokeless tobacco: The folklore and social history of snuffing, sneezing, dipping, and chewing. *Journal of the American Dental Association*, 105, 821–829.
- Cochran, W. G. (1997). *Sampling techniques* (3rd ed.). New York, NY: Wiley.
- Collier, D. J., & Beales, I. J. P. (1989). Drinking among medical students: A questionnaire survey. *British Medical Journal*, 299, 19–22.
- Croen, L. G., Woesner, M., Herman, M., & Reichgott, M. (1997). A longitudinal study of substance use and abuse in a single class of medical students. *Academic Medicine*, 72, 376–381.
- Crofton, J. W., Freour, P. P., & Tessier, J. F. (1994). Medical education on tobacco: Implications of a worldwide survey. *Medical Education*, 28, 187–196.
- Dania, M. G., Ozoh, O. B., & Bande, E. O. (2015). Smoking habits, awareness of risks, and attitude toward tobacco control policies among medical students in Lagos, Nigeria. *Annals of African Medicine*, 14, 1–7.
- Dekker, H. M., Looman, C. W., Adriaanse, H. P., & van der Maas, P. J. (2008). Prevalence of smoking in physicians and medical students, and the generation effect in the Netherlands. *Social Science & Medicine*, 36, 817–822.
- Department of Health. (1995). *Sensible drinking: The report of an inter-departmental working group*. London: HMSO.
- Desalu, K. O., Isheh, K. R., Olokoba, A. B., Salawu, F. K., & Danburam, A. (2010). Smokeless tobacco use in an adult Nigerian population. *Nigerian Journal of Clinical Practice*, 13(4), xx–xx.
- Desalu, O. O., Adekoya, A. O., Elegbede, A. O., Dosunmu, A., & Kolawole, T. F. (2009). Knowledge of and practices related to smoking cessation among physicians in Nigeria. *Journal of the Brazilian Pneumology Society*, 35(11), 1198–1203.
- Desalu, O. O., Olokoba, A. B., Danburam, A., Salawu, F., & Issa, B. M. (2008). Epidemiology of tobacco smoking among the adult population in northeast Nigeria. *Internet Journal of Epidemiology*, 6(1), xx–xx.

- Di Pranza, J. (2010). A new approach to the diagnosis of tobacco addiction. *Addiction*, 105(3), 381–382.
- DiClemente, C. C., & Prochaska, J. O. (1985). Processes and stages of self-change: Coping and competence in smoking behavior change. In S. Schiffman (Ed.), *Coping and substance use* (pp. 319–343). New York, NY: Academic Press.
- Diethelm, P., & McKee, M. (2006). Lifting the smoke screen: Tobacco industry strategy to defeat smoke-free policies and legislation. *European Respiratory Society and Institut du Cancer*.
- Drinking, smoking, and illicit drug use among 15- and 16-year-olds in the United Kingdom. (1996). *British Medical Journal*, 313, 394–397.
- Dumitrescu, A. L. (2007). Tobacco and alcohol use among Romanian dental and medical students: A cross-sectional questionnaire survey. *Oral Health & Preventive Dentistry*, 5, 279–284.
- Edens, E. L., Glowinski, A. L., Pergadia, M. L., Lessov-Schlaggar, C. N., & Bucholz, K. K. (2010). Nicotine addiction in light-smoking African American mothers. *Journal of Addiction Medicine*, 4, 55–60.
- Eysenck, H. J., & Eysenck, S. B. G. (1975). *Manual of the Eysenck Personality Questionnaire (Junior and Adult)*. London: Hodder & Stoughton.
- Ezzati, M., Lopez, A. D., Rodgers, A., Vander, H. S., & Murray, C. J. (2002). Selected major risk factors and global and regional burden of disease. *The Lancet*, 360, 1347–1360.
- Fadhil, I. (2009). Tobacco education in medical schools: A survey among primary care physicians in Bahrain. *Eastern Mediterranean Health Journal*, 15, 969–975.
- Faggiano, F., Versino, E., & Lemma, P. (2001). Decennial trends of social differentials in smoking habits in Italy. *Cancer Causes & Control*, 12, 665–671.
- Faseru, B., Barengo, N. C., Sandström, H. P., & Omokhodion, F. (2006). Smoking behavior and perception of risk among medical students in Ibadan, Nigeria. *Preventive Control*, 2, 103–109.
- Fernandez, E., Schiaffino, A., & Garcia, M. (2001). Widening social inequalities in smoking cessation. *Journal of Epidemiology & Community Health*, 55, 729–730.
- First Annual Report and National Plan. (1999). The United Kingdom Anti-Drugs Coordinator. Central Office of Information, London.
- Fischer, S., Spiegelhalter, B., Eisenbarth, J., & Preussman, R. (1990). Investigations on the origin of tobacco-specific nitrosamines in the mainstream smoke of cigarettes. *Carcinogenesis*, 11(5), 723–730.
- Forney, M. A., Ripley, W. K., & Forney, P. D. (1988). A profile and prediction study of problem drinking among first-year medical students. *International Journal of the Addictions*, 23, 767–779.
- Fowles, J., & Dybing, E. (2003). Application of toxicological risk assessment principles to the chemical constituents of cigarette smoke. *Tobacco Control*, 12, 424–430.
- Frank, E., Denniston, M., & Pederson, L. (2002). Declines in smokers' understanding of tobacco hazards between 1986 and 1998: A report from North Georgia. *Southern Medical Journal*, 95, 675–680.
- Freeman, M. K., Fleming, T. D., Robinson, M., Dwyer-Lindgren, L., Thomson, B., Wollum, A., et al. (2014). Smoking prevalence and cigarette consumption in 187 countries: Report on 1980–2012. *Journal of Public Health Reports*, 129(1), 44–54.
- Gallaway, M. S., Glover-Kudon, R., Momin, B., Puckett, M., Lunsford, N. B., Ragan, K. R., et al. (2019). Smoking cessation attitudes and practices among cancer survivors—United States, 2015. *Journal of Cancer Survivorship*, 13, 66–74.
- Geller, A. C., Brooks, D. R., Powers, C. A., Brooks, K. R., Rigotti, N. A., & Bogner, B. (2008). Tobacco cessation and prevention practices reported by second and fourth-year students at US medical schools. *Journal of General Internal Medicine*, 23, 1071–1076.
- Ghodse, A. H., & Howse, K. (1994). Substance use among medical students: A nationwide survey. *Health Trends*, 26, 85–88.
- Gilman, S. E., Abrams, D. B., & Buka, S. L. (2003). Socioeconomic status over the life course and stages of cigarette use: Initiation, regular use, and cessation. *Journal of Epidemiology & Community Health*, 57, 802–808.
- GlaxoSmithKline. (2009). *Prescribing information*.
- Global Adult Tobacco Survey (GATS). (2012). Nigeria Executive Summary. *Global Facts Sheets*.
- Global Youth Tobacco Survey (GYTS). (2006). Report on the results of GYTS Kenya.

- Retrieved from <http://www.cdc.gov/tobacco/global/GYTS/reports/afro2006/Kenya-report.html>.
- Goldberg, D., & Williams, P. (1988). *Users guide to the General Health Questionnaire*. Windsor, UK: NFER.
- Janz, N., & Becker, M. H. (1984). The health belief model: A decade later. *Health Education Quarterly*, 11, 1–47.
- Jarvis, M. J., & Wardle, J. (1999). Social patterning of individual health behaviours: The case of cigarette smoking. In M. Marmot & R. G. Wilkinson (Eds.), *Social determinants of health* (pp. xx–xx). Oxford: Oxford University Press.
- Royal College of Physicians. (Year). Alcohol and the heart in perspective: Sensible limits reaffirmed. Summary of the report of a working group of the Royal College of Physicians, Psychiatrists, and General Practitioners. *Journal of the Royal College of Physicians London*, 29, 266–271.
- Weiss, Y., Bristow, B., Karol, D. L., Fitch, M., McAndrew, A., Gibson, L., et al. (2020). Exploring tobacco use and smoking cessation best practices from the perspectives of individuals with lung cancer and healthcare professionals. *Journal of Medical Imaging and Radiation Sciences*, 51(1), 62–67.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/127866>