



# Quality of Bread Wheat (*Triticum aestivum* L.) as Influenced by Different Nutrient Management Approaches under Northern Transitional Zone of Karnataka

Ummesalma Sanadi <sup>a\*</sup>, K. K. Math <sup>a</sup>, Kumar D. Lamani <sup>b</sup>  
and Uday G. Reddy <sup>c</sup>

<sup>a</sup> Department of Soil Science and Agricultural Chemistry, University of Agricultural Sciences, Dharwad- 580 005, Karnataka, India.

<sup>b</sup> Department of Agronomy, University of Agricultural Sciences, Dharwad- 580 005, Karnataka, India.

<sup>c</sup> Department of Genetics and Plant Breeding, University of Agricultural Sciences, Dharwad- 580 005, Karnataka, India.

## Authors' contributions

This work was carried out in collaboration among all authors. Author US designed the study, performed the statistical analysis, wrote the protocol, wrote the first draft of the manuscript, managed the analyses. Authors KKM, KDL and UGR managed the literature searches and corrected the manuscript. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJPSS/2023/v35i12711

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/95927>

Original Research Article

Received: 19/11/2022

Accepted: 25/01/2023

Published: 26/01/2023

## ABSTRACT

The field experiments were conducted at the Research Farm of AICW&BIP, Main Agricultural Research Station, University of Agricultural Sciences, Dharwad during *rabi* 2020-21 and 2021-22 and was laid out in Randomized Complete Block Design with 15 treatments replicated thrice to

\*Corresponding author: E-mail: ummesalmarsanadi@gmail.com;

evaluate the performance of wheat quality to various nutrient management approaches. The pooled data onto two years indicated that application of 275:40.33:135.2 kg of N:P<sub>2</sub>O<sub>5</sub>:K<sub>2</sub>O ha<sup>-1</sup> for yield target of 55 q ha<sup>-1</sup> under SSNM practices significantly increased the protein content (13.43 %), wet gluten (34.7 %), dry gluten (11.9 %), gluten index (81.6 %), sedimentation value (46.3 ml) and yellow pigment (5.93 ppm) in wheat grain and it was found on par with yield targets at 50 and 45 q ha<sup>-1</sup> under SSNM.

**Keywords:** Quality; SSNM; STCR; STL; targeted yield; wheat.

## 1. INTRODUCTION

Wheat (*Triticum aestivum* L.) is popularly known as the king of cereals. It is the second most important cereal crop after rice in India and it belongs to family *poaceae*. Wheat is an important source of carbohydrates. Globally, it is the leading source of vegetable protein in human food, having a protein content of about 13 per cent, which is relatively high compared to other major cereals but relatively low in protein quality for supplying essential amino acids, but it is a good source of multiple nutrients (i.e. Phosphorous, Magnesium, Niacin, Iron and Calcium) and dietary fiber. It provides a balanced nutrition to millions of people.

“Wheat ranks first in the area and production at global level, and India is the second largest wheat producer of the World followed by China. In India, the wheat is grown over an area of 31.35 million hectares with annual production of 107.86 million tones and productivity of 3440 kg ha<sup>-1</sup> during 2020-21” [1]. “In Karnataka, wheat is grown over an area of 1.50 million hectares with the production of 1.63 million tones and productivity of 1198 kg ha<sup>-1</sup>” [1].

In India wheat is ground to prepare flour which is mainly consumed after preparing leavened bread (chapati). Its flour is also used to prepare fried chapatti called ‘puris’ and ‘paratha’. In addition to this, wheat is also consumed in various other preparations such as ‘dalia’, ‘halwa’, ‘sweat meal’ etc. One of the most important uses of wheat is to manufacture flour to prepare bake bread, pastry, biscuits etc.

“Nowadays, farmers are trying to get high grain yields in line with food quality, at the same time trying to minimize production costs and to use environmental friendly technologies. Nowadays, not only yield but also the quality of the produced grain is important, because the quality of the grains determines their direction of use” [2]. That is why farmers are trying to get high grain yields in line with food (accepted for bread baking)

quality, while minimize production costs and using environmentally-friendly technologies. Among them balanced nutrition plays an important role in physiological and biochemical processes of plant in determining the yield as well as the quality of produce. In this regard, Soil Test Laboratory (STL), Site Specific Nutrient Management (SSNM), Soil Test Crop Response (STCR) and Nutrient Expert (NE) approaches are unique in the sense that these methods not only indicate soil test based fertilizer dose but also the level of yield the farmer can hope to achieve if good nutrient management practices is followed in raising the crop. Hence, the present investigation are proposed to the objective to evaluate the effect of different fertilizer recommendation approaches on the quality of irrigated wheat.

## 2. MATERIALS AND METHODS

The field experiments were conducted at Research Farm of All India Co-ordinated Wheat and Barley Improvement Project (AICW&BIP), Main Agricultural Research Station, University of Agricultural Sciences, Dharwad during *rabi* 2020-21 and 2021-22 under *irrigated* condition. The study area was located in Northern Transition Zone (Zone VIII) of Karnataka and is situated at 15° 26' North latitude, 75° 07' East longitude and at an altitude of 678 m above mean sea level (MSL). The mean annual rainfall for the past 70 years (1950-2020) was 850.1 mm. The Maximum rainfall of 323.6 mm was received in the month of August followed by October (202.0 mm) during 2020-21. Similarly, during 2021-22 maximum rainfall received during 2021-22 was 187.4 mm in the month of July followed by November (156.2 mm). The mean monthly maximum temperatures of 35.8 °C and 35.4 °C were recorded in April and minimum temperatures of 14.6°C and 13.2 °C were recorded in January, respectively during 2020-21 and 2021-22. The maximum relative humidity of 89 and 87 per cent was recorded in the month of August during 2020-21 and 2021-22, respectively. The wheat variety used for the experiment was UAS-334 (Bread wheat).

**Table 1. Initial soil properties**

<b>Sl. No.</b>	<b>Particulars</b>	<b>Values</b>	<b>Method</b>
1.	Textural class	Clay loam	
2.	Soil reaction (1:2.5, soil water suspension)	7.67	Potentiometric method
3.	Electrical conductivity (1:2.5, soil water extract) (dS m <sup>-1</sup> )	0.26	Conductometric method
4.	Organic carbon (%)	0.68	Walkley and Black's wet oxidation method
5.	Free calcium carbonate (g kg <sup>-1</sup> )	5.42	Rapid acid neutralization method
6.	Available N (kg ha <sup>-1</sup> )	176.9	Modified alkaline permanganate method
7.	Available P <sub>2</sub> O <sub>5</sub> (kg ha <sup>-1</sup> )	33.77	Olsen's method of extraction followed by Spectrophotometric method
8.	Available K <sub>2</sub> O (kg ha <sup>-1</sup> )	361.68	Neutral Normal Ammonium acetate extraction followed by Flame photometric method
9.	Available S (kg ha <sup>-1</sup> )	25.63	0.15% CaCl <sub>2</sub> . 2 H <sub>2</sub> O extraction followed by Turbidimetry
10.	DTPA - extractable micronutrients (mg kg <sup>-1</sup> )		DTPA extraction followed by Atomic absorption spectrophotometric method
a.	Copper	0.58	
b.	Iron	4.34	
c.	Manganese	6.18	
d.	Zinc	0.51	

**Table 2. Treatment details and quantity of fertilizers applied**

Treatment details		Quantity of fertilizers applied (kg ha <sup>-1</sup> )		
		N	P <sub>2</sub> O <sub>5</sub>	K <sub>2</sub> O
T <sub>1</sub>	Absolute control	-	-	-
T <sub>2</sub>	Recommended Dose of Fertilizer (RDF)	100	75	50
T <sub>3</sub>	Recommended Package of Practice (RPP)	100	75	50
T <sub>4</sub>	125% Recommended Dose of Fertilizer (RDF)	125	93.75	62.5
T <sub>5</sub>	150% Recommended Dose of Fertilizer (RDF)	150	112.5	75
T <sub>6</sub>	Soil Test Laboratory (STL) based NPK application	125	75	25
T <sub>7</sub>	SSNM yield target @ 40 q ha <sup>-1</sup>	200.4	29.33	98.4
T <sub>8</sub>	SSNM yield target @ 45 q ha <sup>-1</sup>	225	33	110.4
T <sub>9</sub>	SSNM yield target @ 50 q ha <sup>-1</sup>	250	36.67	123.2
T <sub>10</sub>	SSNM yield target @ 55 q ha <sup>-1</sup>	275	40.33	135.2
T <sub>11</sub>	STCR yield target @ 40 q ha <sup>-1</sup>	170.7	0.0	20.04
T <sub>12</sub>	STCR yield target @ 45 q ha <sup>-1</sup>	208.4	0.0	32.49
T <sub>13</sub>	STCR yield target @ 50 q ha <sup>-1</sup>	246.1	0.0	44.94
T <sub>14</sub>	STCR yield target @ 55 q ha <sup>-1</sup>	283.8	7.25	57.39
T <sub>15</sub>	Nutrient Expert yield target @ 40 q ha <sup>-1</sup>	80	53	45

Before sowing, wheat seeds were treated with Azospirillum @ 3 kg ha<sup>-1</sup>. Seeds were sown in net plot area of 2.0 m x 5.0 m (gross plot: 2.4 m x 6.0 m) at a distance of 30 cm in furrows placed at 5 cm apart. The crop duration was 4 months during first season (29/11/2020 to 24/03/2021) and second season (13/11/2021 to 18/03/2022). It is grown in *irrigated* condition i.e. the crop has given 9 irrigations during its growth. The crop was harvested at physiological maturity when the plants turned yellow with dry straw and grains become hard and contained 20 to 25 per cent moisture. The initial soil properties are presented in Table 1.

The chemical fertilizers were applied as per treatments. Recommended nitrogen, phosphorus and potassium were applied through urea, single super phosphate (16.0 % P<sub>2</sub>O<sub>5</sub>) and muriate of potash (60.0 % K<sub>2</sub>O), respectively. Recommended Fe and Zn were applied in the form of iron sulphate and zinc sulphate. The entire quantity of fertilizer mixture containing entire dose of phosphorus, potassium, iron sulphate and zinc sulphate were applied to each plot at the time of sowing. Crude protein content of wheat grain was determined by multiplying nitrogen concentration in grain by 5.70 and expressed on per cent. The sedimentation values and yellow pigment in wheat grain was analyzed by following standard analytical procedure as described by Mishra and Gupta [3]. Gluten parameters were determined according to standard AACC methods. It was done by using gluten washers (Erkaya, GW 2200<sup>®</sup>, Ankara, Turkey). Ten gram wheat flour samples were analyzed for wet gluten (WG), dry gluten (DG) and gluten index (GI). The following formula was used to calculate the gluten index.

$$\text{Gluten index (GI)} = \frac{\text{Wet gluten remained on sieve (g)}}{\text{Total wet gluten (g)}} \times 100$$

### 3. RESULTS AND DISCUSSION

#### 3.1 Crude Protein Content

“Among the quality traits, grain protein content of wheat has received special attention as a conventional indicator for measuring the nutritional value of food” [4]. Among the different treatments, T<sub>10</sub>: yield target of 55 q ha<sup>-1</sup> under SSNM approaches recorded higher protein content (13.43 %) in wheat grain and was statistically on a par with yield targets of 50 (T<sub>9</sub>) (13.29 %) and 45 q ha<sup>-1</sup> (T<sub>8</sub>) (13.02 %) under the

said practice but these treatments were significantly superior to the remaining treatments (Table 3 and Fig. 1). The crude protein content of wheat grain of individual years ranged from 9.84 (T<sub>1</sub>) to 13.17 per cent (T<sub>10</sub>) in 2020-21 and 9.60 (T<sub>1</sub>) to 13.70 per cent (T<sub>10</sub>) in 2021-22 and the effect of different fertilizer recommendation practices on crude protein content in wheat grain was similar to pooled analysis. The increase of the grain protein content of wheat grain of the said treatments might be due to more nitrogen accumulation in the plants [5]. Nitrogen being the precursor of protein increased its content in grain [6]. Sunil et al. [7] also found that protein content to wheat grain was significantly higher with SSNM approach to fertilizer recommendation to crops in rice-wheat cropping sequence followed by STCR approach of fertilizer recommendation and graded levels of fertilizer application. Increasing levels of nitrogen application significantly increased the grain protein content of wheat [8]. “The increase in protein contents of wheat grain to higher nitrogen application rates might be attributed to the higher N supply that favors the conversion of carbohydrates into proteins, which in turn promotes the formation of protoplasm resulting in translocation of more nitrogen to grains and ultimately increasing the grain protein content” [9]. The results were in conformity with Majid et al. [10], Haile et al. [11] and Youssef et al. [12] in wheat.

#### 3.2 Gluten Content

“According to the solubility of protein components in different solvents, wheat protein can be divided into gliadin, glutenin, albumin and globulin” [13]. “Among them, gliadin and glutenin are the main storage proteins of wheat, and the main constituents of wet gluten. The wet and dry gluten contents and gluten index composition affect the viscoelasticity and baking quality of wheat dough” [14]. Water retention capacity, loaf volume and dough strength of bread are functions of gluten.

The higher wet (34.7 %) and dry (12.3 %) gluten contents and gluten index (81.6 %) of wheat grain was observed in target yield of 55 q ha<sup>-1</sup> (T<sub>10</sub>) and was on a par with yield targets at 50 (T<sub>9</sub>) and 45 q ha<sup>-1</sup> (T<sub>8</sub>) the same practice of nutrient management but significantly superior to rest of the treatments (Table 4 and Fig. 2). The wet gluten content of wheat flours during individual years ranged from 25.4 (T<sub>1</sub>) to 34.0 per cent (T<sub>10</sub>) in 2020-21 and 24.8 (T<sub>1</sub>) to




Fig. 1. Crude protein content as influenced by various nutrient management approaches

Table 3. Effect of different fertilizer recommendation approaches on wheat quality

Treatments	Protein content (%)			Sedimentation value (ml)			Yellow pigment (ppm)		
	2020	2021	Pooled	2020	2021	Pooled	2020	2021	Pooled
T <sub>1</sub> : Absolute control	9.84	9.60	<b>9.72</b>	28.3	27.9	<b>28.1</b>	3.70	3.58	<b>3.64</b>
T <sub>2</sub> : RDF	11.88	11.21	<b>11.54</b>	36.4	36.7	<b>36.6</b>	4.37	4.50	<b>4.43</b>
T <sub>3</sub> : RPP	11.97	12.26	<b>12.11</b>	37.5	37.9	<b>37.7</b>	4.53	4.66	<b>4.60</b>
T <sub>4</sub> : 125% RDF	12.10	12.16	<b>12.13</b>	39.1	39.5	<b>39.3</b>	5.17	5.30	<b>5.23</b>
T <sub>5</sub> : 150% RDF	12.41	12.83	<b>12.61</b>	39.2	39.5	<b>39.4</b>	5.27	5.40	<b>5.33</b>
T <sub>6</sub> : STL approach	11.97	12.43	<b>12.20</b>	38.6	39.3	<b>38.9</b>	4.60	4.70	<b>4.65</b>
T <sub>7</sub> : SSNM yield target at 40 q ha <sup>-1</sup>	12.54	12.84	<b>12.69</b>	41.3	41.7	<b>41.5</b>	5.57	5.70	<b>5.63</b>
T <sub>8</sub> : SSNM yield target at 45 q ha <sup>-1</sup>	12.83	13.22	<b>13.02</b>	42.6	43.0	<b>42.8</b>	5.67	5.80	<b>5.73</b>
T <sub>9</sub> : SSNM yield target at 50 q ha <sup>-1</sup>	13.01	13.57	<b>13.29</b>	44.5	44.8	<b>44.7</b>	5.73	5.83	<b>5.78</b>
T <sub>10</sub> : SSNM yield target at 55 q ha <sup>-1</sup>	13.17	13.70	<b>13.43</b>	46.1	46.5	<b>46.3</b>	5.90	5.97	<b>5.93</b>
T <sub>11</sub> : STCR yield target at 40 q ha <sup>-1</sup>	11.97	12.43	<b>12.21</b>	35.4	36.2	<b>35.8</b>	4.13	4.16	<b>4.15</b>
T <sub>12</sub> : STCR yield target at 45 q ha <sup>-1</sup>	11.38	11.54	<b>11.46</b>	35.8	38.2	<b>37.0</b>	4.57	4.73	<b>4.65</b>
T <sub>13</sub> : STCR yield target at 50 q ha <sup>-1</sup>	12.12	12.84	<b>12.48</b>	39.1	39.0	<b>39.0</b>	5.13	5.26	<b>5.20</b>
T <sub>14</sub> : STCR yield target at 55 q ha <sup>-1</sup>	12.56	13.11	<b>12.83</b>	40.1	40.4	<b>40.2</b>	5.43	5.56	<b>5.50</b>
T <sub>15</sub> : NE yield target at 40 q ha <sup>-1</sup>	11.78	12.08	<b>11.93</b>	35.4	35.7	<b>35.5</b>	4.03	4.26	<b>4.15</b>
SEm±	0.23	0.24	<b>0.16</b>	0.84	0.88	<b>0.73</b>	0.10	0.08	<b>0.08</b>
CD(0.05)	0.65	0.69	<b>0.45</b>	2.43	2.54	<b>2.12</b>	0.29	0.22	<b>0.23</b>
CV (%)	3.2	3.4	<b>2.2</b>	3.8	3.9	<b>3.3</b>	3.5	2.7	<b>2.8</b>

Table 4. Effect of different fertilizer recommendation approaches on wheat quality

Treatments	Wet gluten (%)			Dry gluten (%)			Gluten index (%)		
	2020	2021	Pooled	2020	2021	Pooled	2020	2021	Pooled
T <sub>1</sub> : Absolute control	25.4	24.8	<b>25.1</b>	7.7	7.5	<b>7.6</b>	69.9	69.3	<b>69.6</b>
T <sub>2</sub> : RDF	30.6	28.9	<b>29.8</b>	9.7	9.0	<b>9.4</b>	73.1	73.3	<b>73.2</b>
T <sub>3</sub> : RPP	30.9	31.6	<b>31.3</b>	9.8	10.1	<b>9.9</b>	73.3	73.5	<b>73.4</b>
T <sub>4</sub> : 125% RDF	31.2	31.4	<b>31.3</b>	9.9	10.0	<b>10.0</b>	73.5	73.9	<b>73.7</b>
T <sub>5</sub> : 150% RDF	32.0	33.1	<b>32.5</b>	10.8	11.2	<b>11.0</b>	75.6	76.0	<b>75.8</b>
T <sub>6</sub> : STL approach	30.9	32.1	<b>31.5</b>	9.8	10.2	<b>10.0</b>	74.3	74.7	<b>74.5</b>
T <sub>7</sub> : SSNM yield target at 40 q ha <sup>-1</sup>	32.4	33.1	<b>32.7</b>	10.9	11.2	<b>11.1</b>	75.9	76.8	<b>76.4</b>
T <sub>8</sub> : SSNM yield target at 45 q ha <sup>-1</sup>	33.1	34.1	<b>33.6</b>	11.2	11.6	<b>11.5</b>	77.5	78.4	<b>78.0</b>
T <sub>9</sub> : SSNM yield target at 50 q ha <sup>-1</sup>	33.6	35.0	<b>34.3</b>	11.4	11.9	<b>11.7</b>	78.1	79.0	<b>78.6</b>
T <sub>10</sub> : SSNM yield target at 55 q ha <sup>-1</sup>	34.0	35.3	<b>34.7</b>	11.5	12.3	<b>11.9</b>	81.2	82.1	<b>81.6</b>
T <sub>11</sub> : STCR yield target at 40 q ha <sup>-1</sup>	29.4	32.1	<b>30.7</b>	9.8	10.2	<b>10.0</b>	72.7	73.2	<b>73.0</b>
T <sub>12</sub> : STCR yield target at 45 q ha <sup>-1</sup>	30.9	29.8	<b>30.4</b>	9.2	9.4	<b>9.3</b>	74.7	75.2	<b>75.0</b>
T <sub>13</sub> : STCR yield target at 50 q ha <sup>-1</sup>	31.3	33.1	<b>32.2</b>	9.9	10.7	<b>10.3</b>	75.6	76.1	<b>75.9</b>
T <sub>14</sub> : STCR yield target at 55 q ha <sup>-1</sup>	32.4	33.7	<b>33.1</b>	10.9	11.5	<b>11.2</b>	76.5	77.0	<b>76.7</b>
T <sub>15</sub> : NE yield target at 40 q ha <sup>-1</sup>	30.4	31.2	<b>30.8</b>	9.6	9.9	<b>9.8</b>	70.7	70.9	<b>70.8</b>
SEm±	0.5	0.5	<b>0.4</b>	0.2	0.2	<b>0.1</b>	1.4	1.4	<b>1.4</b>
CD(0.05)	1.4	1.5	<b>1.2</b>	0.5	0.7	<b>0.4</b>	4.0	4.0	<b>4.0</b>
CV (%)	3.3	3.4	<b>2.2</b>	3.0	4.0	<b>2.6</b>	3.3	3.2	<b>3.2</b>







Fig. 2. Gluten content of wheat as influenced by various nutrient management approaches



**Fig. 3. Sedimentation value of wheat as influenced by various nutrient management approaches**

35.3 per cent ( $T_{10}$ ) in 2021-22. Similarly, dry gluten content of wheat flour was similar in both the individual years and the value ranged from 7.7 ( $T_1$ ) to 11.5 per cent ( $T_{10}$ ) during 2020-21 and 7.5 ( $T_1$ ) to 12.3 per cent ( $T_{10}$ ) during 2021-22. The gluten indexes of wheat grain of individual years ranged from 69.9 ( $T_1$ ) to 81.2 per cent ( $T_{10}$ ) in 2020-21 and 69.3 ( $T_1$ ) to 82.1 per cent ( $T_{10}$ ) in 2021-22. This is because of higher protein content of wheat grain in the above said treatments, as there is synergetic relation between wheat grain protein and gluten contents. This is also attributed to application of nitrogen in adequate quantity as per the crop requirement which might have improved the quality of grains because nitrogen is an essential component of protein. The present results can be correlated with the findings of Litke et al. [15] who opined that application of  $240 \text{ kg N ha}^{-1}$  to wheat grown in loam soils increased the protein and gluten contents from 8.7 to 13.7 per cent and 14.73 to 28.74 per cent, respectively.

### 3.3 Sedimentation Value


Sedimentation values of wheat was significantly higher in  $T_{10}$ : SSNM yield targeted at  $55 \text{ q ha}^{-1}$  (46.3 ml) and was on a par with  $50 \text{ q ha}^{-1}$  ( $T_9$ )(44.7 ml) under the same nutrient management practice and both of these treatments were significantly superior over rest of the treatments (Table 3 and Fig. 3). The yield target of 40 and  $45 \text{ q ha}^{-1}$  under SSNM approach recorded statistically similar sedimentation value.

The impact on various fertilizer recommendation approaches on sedimentation value of wheat flour was similar in both the individual years and the values ranged from 28.3 ( $T_1$ ) to 46.1 ml ( $T_{10}$ ) during 2020-21 and 27.9 ( $T_1$ ) to 46.5 ml ( $T_{10}$ ) during 2021-22. This is because sedimentation values is a gluten dependent quality parameter and is directly related to gluten content in wheat grain. This is based on the fact that gluten protein absorbs water and swells considerably when treated with lactic acid in the presence of sodium dodecyle sulphate (SDS).

The higher sedimentation value of wheat flour might be due to higher expansion and sedimentation of gluten that led to high-quality and high-strength gluten [16]. Therefore, sedimentation value can be used as an essential indicator for detecting the quality of gluten.

### 3.4 Yellow Pigment

Yellow pigment content is another quality parameter that imparts attractive yellow colour to the wheat products. It is precursor of vitamin 'A' and hence has immense nutritional importance. Higher yellow pigment in wheat grain was recorded in the treatment for target yield at  $55 \text{ q ha}^{-1}$  (5.93 ppm) under SSNM practice ( $T_{10}$ ) and was on a par a with yield targets of  $50 \text{ q ha}^{-1}$  ( $T_9$ )(5.78 ppm) and  $45 \text{ q ha}^{-1}$  ( $T_8$ )(5.73 ppm) under the same nutrient management practices but differed significantly with other treatments (Table 3 and Fig. 4). Yellow pigment in wheat grain of



**Fig. 4. Sedimentation value of wheat as influenced by various nutrient management approaches**

individual years ranged from 3.70 (T<sub>1</sub>) to 5.90 ppm (T<sub>10</sub>) in 2021-22 and 3.58 (T<sub>1</sub>) to 5.97 ppm (T<sub>10</sub>) in 2021-22 and the results were similar to pooled analysis. This is attributed to improvement in grain quality traits such as protein and gluten contents in grain which significantly influenced the yellow pigment.

#### 4. CONCLUSION

Fertilization on the basis of soil test is one of the best and effective ways to improve yield and nutritional quality of crops for human consumption. Among them site specific nutrient management approaches for yield targeted of 55 q ha<sup>-1</sup> is best for getting higher protein content, gluten content, sedimentation value and yellow pigment in bread wheat over other approaches. Protein content of wheat flours is one of the main factors that determine nutritional and commercial quality of wheat, and it is greatly influenced by available nitrogen in the soil. Grain protein content usually increases when plant available nitrogen is not a limiting factor for grain yield. Balanced application of nutrients during crop growth period helps in increasing protein content of wheat flour and also improves the gluten content, sedimentation value and yellow pigment.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Anonymous. Agricultural Statistics. Directorate of Economics and Statistics, Department of Agriculture and Cooperation, Ministry of Agriculture, Government of India, New Delhi; 2021.
2. Jin H, Wen W, Liu J, Zhai S, Zhang Y, Yan J. Genome-wide QTL mapping for wheat processing quality parameters in a gaocheng 8901/ zhoumai 16 recombinant inbred line population. *Front. Plant Sci.* 2016;7:1032.
3. Mishra BK, Gupta RK. Protocols for evaluation of wheat quality. Directorate of Wheat Research, Karnal. 1995;43-48.
4. Zhao LA, Zhang KP, Liu B, Deng ZY, Qu HL, Tian JC, A comparison of grain protein content QTLs and flour protein content QTLs across environments in cultivated wheat. *Euphytica.* 2010; 174(3):325–335.
5. Singh V. Productivity and economics of rice (*Oryza sativa*)-wheat (*Triticum aestivum* L.) cropping system under

- integrated nutrient management supply system in recently reclaimed sodic soil. Indian J. Agron. 2017;51(2):81-84.
6. Dinesh PS, Dashrath S, Effect of nitrogen and FYM on yield, quality and uptake of nutrients in wheat (*Triticum aestivum*). Ann. Plant Sci. 2017;19(2):232- 236.
  7. Sunil K, Panwar AS, Naresh RK, Prem S, Mahajan NC, Udit C, Sudhir K, Minaxi M, Meena AL, Ghashal PC, Meena LK, Jairam C. Improving rice-wheat cropping system through precision nitrogen management: A review. J. Pharm. Phytochem. 2018;7(2):1119-1128.
  8. Halverson AD, Nielsen DC, Reule CA. Nitrogen Fertilization and rotation effects on no-till dry land wheat production. Agronomy. 2004;96:1196-1201.
  9. Zemichael B, Dechassa N, Fetien. Yield and nutrient use efficiency of bread wheat (*Triticum aestivum* L.) as influenced by time and rate of nitrogen application in Enderta, Tigray, Northern Ethiopia. Open Agriculture. 2017;2:611–624.
  10. Majid MA, Wyseure GCL, Biswas SK, Hossain ABMZ. Farmer's perceptions and knowledge in using wastewater for irrigation in wheat at twelve peri-urban areas and two sugar mill areas in Bangladesh. Agric. Water Manag. 2010;98:79-86.
  11. Haile D, Nigussie D, Ayana A. Nitrogen use efficiency of bread wheat: Effects of nitrogen rate and time of application. J. Soil Sci. Plant Nutr. 2012;12(3):389-409.
  12. Youssef SM, Faizy S, Mashali SA, El-Ramady, Ragab. Effect of different levels of NPK on wheat crop in North Delta. Int. J. Biosen. Bioelectron. 2013;7(12):501-506.
  13. Singh J, Skerritt JH. Chromosomal control of albumins and globulins in wheat grain assessed using different fractionation procedures. J. Cereal Sci. 2001;33(2):163-181.
  14. Torbica A, Antov M, Mastilovic J, Knezevic D. The influence of changes in gluten complex structure on technological quality of wheat (*Triticum aestivum* L.). Int. Food Res. J. 2007;40(8): 1038–1045.
  15. Litke L, Gaile Z, Ruza A. Effect of nitrogen fertilization on winter wheat yield and yield quality. Agronomy. 2018;16(2):500-509.
  16. Pena RJ. Wheat for bread and other foods in Bread wheat improvement and production. Euphytica. 2002;30:483–494.

© 2023 Sanadi et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/95927>